

PIGLET MORTALITY IN DANISH ORGANIC HERDS

Lena Rangstrup-Christensen, DVM, PhD Fellow

Department of Animal Science, Aarhus University, Denmark

Supervisors:

Jan Tind Sørensen & Lene Juul Pedersen

THE PROJECT

- ▶ **Why?** An estimated preweaning mortality between 25 and 40%
- ▶ **What?** Identifying piglet mortality risk factors in Danish organic herds
- ▶ **How?**
 1. Registrations on the piglet mortality during one year
 2. Necropsies on a sample of dead piglets

ON-FARM MORTALITY REGISTRATIONS


- ▶ A pocket sized booklet.
- ▶ Sows: Parity, body condition, lameness etc.
- ▶ Counting and registration of piglets;
 - 1st inspection after birth
 - Castration (or 3-5 days after birth)
 - Vaccination (or 14-21 days after birth)
 - Weaning (7 weeks after birth)
- ▶ June 2014 until May 2015.

Registrations on approximately 6000 farrowings


Hole:	So nr.	590	774	911	635	642	205
	Mytte	73	63	26	115	65	44
	Laug nr	5	2	1	4	4	3
	Fen nr/fælles fold						
Indsæt faremark	Dato	23/6	25/6	25/6	25/6	25/6	25/6
	Marthed (0-1) + Huld (2-4)	3	1	1	3	2	3
	Faring, dato	24/6	24/6	24/6	24/6	24/6	24/6
	L. tilsyn efter faring, dato						
Faring	Dødfødt	5	3		6	2	1
	Dødt efter faring (S/A)		1	5		3	5
	Levendefødt	19	20	3	18	24	22
	Heraf små (max. 21 cm)		2			1	
Indtil kastrering	Kuldudjævning +/-	-1	-5	+9	-3	-6	
	Antal pletter	14	19	19	14	14	15
Gylte	Kastrering dato	30/6	30/6	30/6	30/6	30/6	30/6
	Antal gris v./start	14	14	12	14	12	13
	Altiv. efter optæl, antal						
	Evt. kuldudjævning +/-						
Forglæd/ vaccination	Dato	13/7	13/7	1	13/7	9/7	13/7
	Antal gris	13	13		10	12	13
Fravæning	Dato	24/6	24/6	14/6	24/6	24/6	24/6
	Antal gris frav. soen	12	11	2	10	12	10
Noter (per upløst):		født børst		SUR		altiv?	

*Løsler: 1. Førelser 2. yverbet, 3. ben 4. andet, Anticubus 5. gylte

PREWEANING MORTALITY - SUMMER 2014



STILLBORN PIGLETS – SUMMER 2014




LITTER SIZE - SUMMER 2014

Herds							
Total litter size	15,90	17,62	17,58	14,73	15,71	14,90	15,07
Live born	14,72	15,66	15,53	13,55	14,77	13,83	13,54
Stillborn	1,18	1,96	2,05	1,18	0,94	1,07	1,53


NECROPSIES

- ▶ Dead piglets from approximately 25 sows on each farm.
- ▶ 4 seasonal collection periods.
- ▶ Necropsies performed at the farms.
- ▶ Cause of death, body condition, stomach content etc.



Necropsies on approximately 4000 piglets



NECROPSIES - SUMMER 2014



NECROPSIES - SUMMER 2014



Time of death - Stillborn piglets

SUMMARY PRELIMINARY RESULTS

- ▶ On farm mortality registrations – high mortality
 - › Hot summer in 2014!
- ▶ Necropsies
 - › Crushed piglets represent the largest group of diagnoses
 - › The majority of the crushed piglets had suckled before dying
 - › The majority of stillborn piglets die during the farrowing



PERSPECTIVES

- ▶ Influence of sows parity, body condition and lameness?
- ▶ Enhanced survival if born in a small rather than in large litter?
- ▶ Higher mortality in piglets with many stillborn litter mates?
- ▶ Are small piglets more prone to crushing?
- ▶ Different patterns in mortality depending on season?
- ▶ Influence of management factors such as the use of nursing sows?



THANK YOU FOR YOUR ATTENTION!

