



NIBIO

NORSK INSTITUTT FOR
BIOØKONOMI

NIBIO REPORT

VOL.: 1, NR.: 1, 2015

AGRICULTURAL WATER MANAGEMENT IN THE NORDIC COUNTRIES

Background document for NJF-seminar 487



FORFATTERE

[Senter eller enhet]

TITTEL/TITLE

AGRICULTURAL WATER MANAGEMENT IN THE NORDIC COUNTRIES

FORFATTER(E)/AUTHOR(S)

M. Bechmann, D. Collentine, Flemming Gertz, Morten Graversgaard , B. Hasler, J. Helin, B. Jacobsen, K. Rankinen, and Karen Refsgaard

| | | | | |
|-------------------|--|-------------------------------|--------------------------------|---------------------------------------|
| DATE/DATE: | RAPPORT NR./ REPORT NO.: | TILGJENGELIGHET/AVAILABILITY: | PROSJEKT NR./PROJECT NO.: | SAKSNR./ARCHIVE NO.: |
| 22.12.2015 | Vol/nr/år | Åpen | Prosjektnr | Arkivnr |
| ISBN-NR./ISBN-NO: | ISBN DIGITAL VERSJON/ ISBN DIGITAL VERSION: | ISSN-NR./ISSN-NO: | ANTALL SIDER/ NO. OF PAGES: | ANTALL VEDLEGG/ NO. OF APPENDICES: |
| ISBN | Versjon nr | ISSN | Antall sider | Vedlegg |

OPPDRAGSGIVER/EMPLOYER:

Oppdragsgiver: Landbruksdirektoratet

KONTAKTPERSON/CONTACT PERSON:

Kontaktperson:

STIKKORD/KEYWORDS:

Stikkord norske

Stikkord engelske

FAGOMRÅDE/FIELD OF WORK:

Sett inn fagområde

Insert field of work

SAMMENDRAG/SUMMARY:

Sammendrag (på norsk obligatorisk hvis åpen rapport på engelsk)

Summary (obligatory if open report in English)

LAND/COUNTRY:

Land

FYLKE/COUNTY:

Fylke

KOMMUNE/MUNICIPALITY:

Kommune

STED/LOKALITET:

Sted

GODKJENT /APPROVED

[Navn Navnesen]

NAVN/NAME

PROSJEKTLEDER /PROJECT LEADER

[Navn Navnesen]

NAVN/NAME



NIBIO

NORSK INSTITUTT FOR
BIOØKONOMI

FORORD

In an NJF-seminar on Agricultural water management researchers, representatives from governmental agricultural agencies at local, regional and national level and farmers' advisory services from the Nordic countries were invited to participate. The aim of the meeting was to better understand the following questions in a comparative way.

- What policies (instruments and measures) have been used over the last decades in the Nordic countries?
- What is the status of knowledge on management, efficiency and effectiveness
- What is successful water management?
 - How can success be measured?
 - What are the results of the policies?
 - What were the drivers for the measures?
 - Could other measures have been used?
 - Bottom-up or Top-down approach: How have the farmers and advisors been included?
 - What are the challenges to not implement measures?
 - Are the policies and the instruments targeted to reach the required abatement?

The present report is a background document for the seminar describing water quality problems, strategies to implement mitigation measures, and costs and effects of mitigation measures in the Nordic countries. The authors from each country as listed below have written the country-specific chapters in the report.

Marianne Bechmann, NIBIO – Norwegian Institute for Bioeconomy, 1430 Ås, Norway

Dennis Collentine, SLU - Swedish University of Agricultural Sciences, Uppsala, Sweden

Flemming Gertz, SEGES, Agro Food Park 15, Skejby, 8200 Aarhus N, Denmark

Morten Graversgaard, Department of Agroecology, Aarhus University, 8830 Tjele, Denmark

Berit Hasler, Department of Environmental Science, Aarhus University, 4000 Roskilde, Denmark

Janne Helin, LUKE, Finland

Brian H. Jacobsen, IFRO, University of Copenhagen, 1958 Frb., Denmark

Katri Rankinen, SYKE, 00251 Helsinki, Finland

Karen Refsgaard, NIBIO – Norwegian Institute for Bioeconomy, 1430 Ås, Norway

Ås, 22.12.15

Marianne Bechmann

INNHold

| | |
|---|--|
| SUMMARY | 6 |
| 1 INTRODUCTION | 9 |
| 2 WATER QUALITY PROBLEMS IN RELATION TO AGRICULTURE | 9 |
| 2.1 Norway | 10 |
| 2.1.1 Inputs of nutrients to coastal waters..... | Fejl! Bogmærke er ikke defineret. |
| 2.1.2 Fresh waters..... | Fejl! Bogmærke er ikke defineret. |
| 2.1.3 Agriculture and eutrophication | Fejl! Bogmærke er ikke defineret. |
| 2.2 Denmark..... | 11 |
| 2.3 Sweden..... | 14 |
| 2.4 Finland..... | 14 |
| 3 IMPLEMENTATION OF MITIGATION MEASURES..... | 16 |
| 3.1 Norway | 16 |
| 3.1.1 Norway's international obligations..... | 16 |
| 3.1.2 Legislation and grant systems..... | 16 |
| 3.1.3 Water protection policies and management in Norway | 17 |
| 3.1.4 Level of implementation and effect..... | 19 |
| Each year Norwegian statistics (SSB) publish time series of statistic information on farmer's behavior and choices in an annual report (Bye et al., 2015). Selected data and text from this report related to nutrient application, soil tillage and specific mitigation methods is presented here below. | |
| 19 | |
| 3.1.5 Norwegian summary..... | 26 |
| 3.2 Denmark | 27 |
| 3.2.1 The Aquatic action plans (1987-2005)..... | 27 |
| 3.2.2 The WFD and the River Basin management Plans from 2011 | 30 |
| 3.2.3 Measures and N-policies 2014 until today | 31 |
| 3.2.4 Danish water councils - a new bottom-up approach to River Basin Management Planning | 31 |
| 3.2.5 Conclusion and discussion - Denmark | 32 |
| 3.3 Sweden..... | 33 |
| 3.3.1 Policy..... | 33 |
| 3.3.2 Measures | 33 |
| 3.3.3 The tax on mineral fertilizers | 34 |
| 3.3.4 Agri-environmental measures supported by the Rural Development Programme (2000-2013) | |
| 34 | |
| 3.3.5 Effect and implementation | 34 |
| 3.3.6 Measure effects and cost effectiveness | 38 |
| 3.3.7 Swedish summary | 40 |
| 3.4 Finland | 40 |

| | | |
|-------|---|----|
| 3.4.1 | Introduction to Water protection policies of agriculture in Finland 1995-2015 | 40 |
| 3.4.2 | Objectives and Coherence | 40 |
| 3.4.3 | Reference level for payments, implementation and compliance..... | 41 |
| 3.4.4 | Evidence base and Policy evaluation | 43 |
| 3.4.5 | Summary - Finland | 45 |
| 4 | CONCLUSION | 47 |
| 5 | LITERATURE | 48 |
| 6 | DANISH APPENDIX : SUMMARY OF CATALOGUE OF MEASURES (ERIKSEN ET AL., 2014)..... | 52 |

UDKAST

SUMMARY

Water quality

The structure of and the conditions for agricultural production in the four Nordic countries, Norway, Denmark, Sweden and Finland, differ with regard to climate and geology, which influences agronomic practices and the political and economic framework. There are large differences in the problems related to water quality. In Norway and Finland, the main focus is on water quality in lakes and streams, and phosphorus (P) has shown to be the limiting nutrient for eutrophication in these water bodies, the focus in these two countries are on reduction in P loading. In Denmark, the main focus is on nitrate in ground water and eutrophication of the fjords and the coastal areas. Therefore, the Danish focus is mainly on nitrogen (N). In Sweden, the problems of eutrophication are both in fresh water and in the Baltic Sea and measures to reduce both N and P have been focused.

Environmental measures

Despite the differences between the countries, the agricultural mitigation measures implemented in the four countries comprise several similar methods. For P, the measures include among others management of manure, changed soil tillage, grassed buffer zones along open water and sedimentation ponds. Measures for reduced N leaching include among others catch crops, constructed wetlands and management of manure. In Denmark, strong restrictions in N application have also been implemented.

Instruments

The four countries have much in common, but there are also large differences between the instrumentation in agricultural policy. In Denmark, general command-and-control-measures are dominating, whereas more enticement is used in Finland and Norway with a high degree of regional adaptations.

In Norway, the implementation of mitigation measures consist of general production grants (among others demanding 2-m buffers and nutrient management plan), legislation on manure management and subsidies for e.g. changed soil tillage, grassed buffers and sedimentation ponds administered through the Regional Environmental Program (RMP) and Special measures in agriculture (SMIL).

The Finnish program for environmental measures is similar to the Norwegian and consists of three different instruments for subsidies, which are both functionally and regionally differentiated. It consists of a basic subsidy, which among others includes maximum N application level and a general claim of good agronomy. In addition, each farm can choose one or two mitigation measures, for example catch crops, reduced nutrient application or manure application in growing season. The third Finnish program comprises specific subsidies which are voluntary for example grassed buffer zones or sedimentation ponds.

In Sweden, implementation of mitigation measures is based on legislations, information campaigns and subsidies. Additionally, tax on N- and P fertilizer has been tested as a mean to reduce nutrient losses, but this is no longer used.

Most of the Danish mitigation measures have been implemented as legislations (Command-and-control). In 2013, a new law on water planning (Lov om vandplanlægning) which emphasize the need for a bottom-up approach (e.g. Danish Water Councils based on the Swedish approach) at the local level was adopted, however the implementation remains.

Norway, Finland and Sweden place much focus on the measures in the Rural Development Programmes (RDP) while most of the Danish agricultural measures are outside this programme. The RDP program is in Denmark mainly used for support to organic farming, new technology and wetlands.

Cost-effectiveness

Targeting of mitigation measures often increases the cost-effectiveness of the measures and in all four countries the focus on targeting mitigation measures has increased lately. On the other hand, targeting often result in high transaction costs in the form of administration, advisory service and control. An evaluation of the balance between targeting of mitigation measures and the transaction costs is still missing. Research shows that knowledge-transfer and involvement in the planning process may result in improved implementation of measures. In each country estimates for cost-effectiveness for agricultural mitigation measures has been calculated.

In Norway, the cost-effectiveness has been evaluated recently by Refsgaard et al. (2013) for changed soil tillage, buffer zones and sedimentation ponds. The most important results were the large variation in cost-effectiveness related to variation in erosion risk, with the best cost-effectiveness obtained by implementing mitigation measures on high risk areas of erosion. As an example, the cost-effectiveness of all soil tillage changes at erosion risk over 2 tons soil/ha were less than 110 €/kg P. Implementation of spring tillage on soil with erosion risk higher than 0.5 kg soil/ha costs less than 220 €/kg P. Mitigations measures in the agricultural sector were shown to be cheap compared to measures in other sectors, but they showed great uncertainty in the effect.

Denmark

Sweden

Finland

Further work in the project

The present report will form the basis of further work on evaluating similarities and differences in cost-effectiveness and the success in instrumentation of agricultural mitigation measures in Denmark, Norway, Finland and Sweden. The work will especially focus on the role of the Rural Development Programmes and the use of subsidies to improve the environment. Is there a clear link between the rural programmes and the environmental aims and what do the economic analysis show regarding level of subsidies and the uptake of the measures? Also the paper will look at the extent to which the measures are targeted and the data used to find the most cost-effective locations. The results of this work will be published in a scientific paper and a Fact sheet from NIBIO.

UDKAST



1 INTRODUCTION

The River Basin Management plans (RBMP) and Programme of Measures (PoM) have been adopted in the EU countries and also in Norway. The plans will be implemented according to the planning cycles in the EU Water Framework Directive (WFD). In most countries there is limited knowledge about the costs of the measures and challenges related to the effectiveness of the instrumentation and the measures. Research shows that legitimacy and knowledge about the costs are important factors for a successful implementation. Further that the costs, effects and farmers adaptation varies with natural and institutional conditions. Therefore better evaluation is needed to calculate cost and effect of the measures as well as analyses of relevant policies to target agricultural pollution.

2 WATER QUALITY PROBLEMS IN RELATION TO AGRICULTURE

2.1 Norway

The status on water quality in Norway is described at the official Norwegian web-page “Environment.no”, which is edited by The Norwegian Environment Agency assigned by The Ministry of the Environment. Selected information is presented here.

Inputs of nutrients to coastal waters are steadily rising in most parts of Norway, whereas inputs along the Skagerrak coast have been reduced since 1985. Measurements show that water quality is generally good in more open areas along the Skagerrak coast, but poorer in certain fjords.

The impacts of nutrient inputs are more marked along the Skagerrak coast than anywhere else in Norway. Agriculture and municipal waste water treatment are the main anthropogenic sources of N and P inputs in this area.

Environmental conditions in Norwegian rivers and lakes are good compared with those in most other countries in Europe. A preliminary survey of the status of all Norwegian water bodies shows that around 50 per cent probably will meet the EU objectives for the freshwater environment, while around a quarter are at risk. The rest lacks data or has uncertain status. There are wide regional variations, and not surprisingly, environmental conditions are poorest where the population density is highest. Despite the introduction of numerous measures in recent years, problems with eutrophication still remain. In the future, climate change is likely to escalate the problems, particularly with regards to increased runoffs.

Norway's rivers and lakes provide a rich and varied freshwater environment, which is under less pressure from human activity than many countries in Europe. Challenges still remain as many rivers and lakes are altered due to hydropower regulation, urbanisation and roads.

Only around 7 per cent of Norway's fresh water is characterised as ground water, and accounts for a mere 15 per cent of the water consumption. This is very low compared to many other countries in Europe and is due to the country's abundant supply of surface water.

Eutrophication is still a problem in many rivers and lakes despite the introduction of numerous measures in recent years. Environmental monitoring shows that the situation has remained unchanged in the last ten years. Climate change is one probable factor causing eutrophication. Most of the eutrophication problems in rivers and lakes are related to P and the measures implemented in Norway are especially focusing on reductions in P loading. The following map shows the effect of P on water quality in rivers and lakes.




Figure 2.1.1. Eutrophication in rivers and lakes in Norway

One source of inputs of nutrients from agriculture is the manure and mineral fertiliser applied to arable land and pasture. N is relatively easily leached directly from the soil, whereas most P is bound to soil particles, but can be washed out into river systems and the sea by soil erosion. Other sources of nutrients are leaks and discharges from silos, slurry storage systems and milking parlours.

Runoff of N and P from farmland causes the most widespread pollution, and can result in persistent eutrophication problems – excessive algal and other plant growth and oxygen depletion in river systems and coastal waters. Discharges from silos and slurry stores can give rise to similar problems, but generally on a more local scale.

Inputs of nutrients from agriculture have remained almost unchanged over the past ten years. Intensification of agriculture tends to result in higher releases of nutrients. Crop yields are often increased by applying more fertiliser per unit area, and more livestock are kept on the same area of land, making it necessary to apply more manure per unit area. In both cases, nutrient runoff will increase.

To avoid this, Norway requires arable farmers to plan fertiliser application to avoid a surplus of nutrients, and there are rules limiting the number of livestock that may be kept per unit area of land. Furthermore, subsidies are given to soil tillage methods that reduce erosion and to retention-measures for nutrients and soil particles (grassed bufferzones and sedimentation ponds).

2.2 Denmark

Denmark has a long history of introducing water quality policies and management relying on a 30 years history of aquatic action plans aiming at reducing nutrient losses from agriculture and discharge from waste water pollution. Eutrophication, caused by nutrient leakage, is regarded as the major water quality problem in Denmark. Most of the plans adopted have focused on reducing N losses from agriculture and the losses have been reduced by 50% from mid 1980'ties until now. Some of the key measures used in this period have been max quotas for N on farm level, improved utilization of nutrients in manure, mandatory catch crops and wetland recreation, and the implementation of these measures have led to a significant reduction in the use of mineral fertilizer and a reduction of more than 50% in the N load to coastal waters. However, the majority of all streams and lakes and almost all coastal waters are below good ecological status so more efficient measures are required to reach this target set in the Water Framework Directive.

Ecological status of surface waters

- Bad
- Good
- Moderate
- High
- Poor





Figure 2.2.1 Ecological status of the Danish surface waters: Streams, Lakes and marine waters (Danish Nature Agency, 2014).

2.3 Sweden

Eutrophication is a problem not only in many inland waters in Sweden but also in the Baltic Sea, where three districts draining to the Baltic Sea are having eutrophication problems. In Sweden each of the five Swedish Water Districts is responsible for ensuring good water status including non-eutrophic status under the European Water Framework Directive (WFD). The Northern Baltic Sea District estimates that around 48% of the water in the district is eutrophic (NBWD, 2008). The Swedish government is also committed as a signatory to the Baltic Sea Action Plan (BSAP) to reduce nutrient loads to the Baltic Sea to achieve good environmental status by 2021 (SNV, 2008). In addition, *No Eutrophication* is one of the 16 environmental quality objectives adopted by the Swedish Parliament. Unfortunately, controlling nutrient losses has been more difficult than anticipated due to the diffuse nature of the loads.

Agriculture is the primary source of diffuse nutrient loads to the aquatic environment. According to a report by the Swedish Environmental Protection Agency (SwEPA, 2009), agricultural practices in 2006 were responsible for 46% of the gross anthropogenic loads of N and P to the environment and around 42% of the net load to the Baltic Sea. Gross anthropogenic N losses from cultivated land in 2005 were estimated to be 49 000 tons and P losses for the same year to be 1 400 tons (Johnsson et al, 2008). While both of these nutrient loss estimates represent improvements over the period from 1995 – 2005 (over which N losses decreased by 14% and P losses by 6,6%) there is still a need for considerable reductions to achieve good water quality status under the Water Framework Directive. The Swedish government commissioned a study to be performed by the Swedish EPA (Naturvårdsverket) and the Swedish Board of Agricultural (Jordbruksverket) to determine how loads of N and P from the agricultural sector could be reduced to meet the targets consistent with the BSAP. The conclusion of this study (SNV, 2008) was that it would not be possible to meet the targets with the set of measures studied.

2.4 Finland

In Finland the recent ecological classification of surface waters showed that rivers and coastal waters need attention in improving their state but larger lakes were mainly in excellent or good state (http://mmm.multiedition.fi/syke/envelope/Envelope_2013_3/sivu_5.php). However, small lakes are suffering from eutrophication. Regional differences in the status of waters are considerable. None of the water areas in coastal regions has high status. The conditions of the Archipelago Sea and of the Gulf of Finland are especially worrying. The status of coastal waters has deteriorated in some parts of the northern region of the Bay of Bothnia and the river mouths of the Gulf of Bothnia.

The most significant problem is the eutrophication of waters. Investments in municipal and industrial waste water purification effectively improved the quality of inland waters (Räike et al. 2003). Nowadays water protection policy concentrates on agriculture as it comprises the largest source of nutrients into water

bodies. Agricultural land use covers only 9% of the total land area but it is concentrated in the southern and western parts of the country. Earlier increased P fluxes had been observed to cause eutrophication of surface waters as P is the major nutrient controlling eutrophication in many aquatic systems. Later studies have shown that the Baltic Sea may be seasonally or spatially N limited (Tamminen and Andersen 2007). Further, P is not always the limiting nutrient in lakes either as some smaller lakes may be also N limited (Pietiläinen and Räike 1999).

The flow normalized total nitrogen (TN) flux to the Baltic Sea increased steadily until the period 2000–2006. Turning point occurred around 2000 in the drainage basins of the Gulf of Finland and the Bothnian Sea, and around 2008 in the drainage basin of the Bothnian Bay. Obvious increase in the flow normalized TN flux occurred from late 1990's to mid 2000's. The flow normalized total phosphorus (TP) flux was highest to the Gulf of Finland and to the Bothnian Bay in 1985–1990. Since then the TP flux steadily decreased, exception is the TP flux to the Archipelago Sea which has remained unchanged in 1985–2012.




Figure 2.4.1 Ecological status of the Finnish surface waters

3 IMPLEMENTATION OF MITIGATION MEASURES

3.1 Norway

3.1.1 Norway's international obligations

Norway has undertaken a number of international obligations to limit or reduce nutrient inputs, for example under the WFD and the North Sea Declarations. The local and central government authorities are coordinating efforts to comply with these obligations (Environment.no).

3.1.2 Legislation and grant systems

At Environment.no the policies involved in implementation of mitigation measures are described as follows. During the 1980s and 1990s, a system of legislation and grants or low-cost loans was developed to encourage farming practices that would reduce diffuse source runoff from agricultural land and point discharges from silos and manure storage systems. The system has been amended and adapted over the years. The legislation includes various regulations under the Pollution Control Act and the Land Act. There are rules on the levelling of steep and hilly farmland to prevent runoff, and regulations on manure and silage effluent that are intended both to reduce point discharges from storage facilities and runoff after application of organic fertilisers.

The regulations relating to production grants set out a number of environmental standards farmers must meet to be entitled to grants, including pesticide journal, fertiliser application plan, and two meter buffer zone along water ways. A farmer who does not comply with the requirements may lose part of the production grant. The Agricultural Directorate is responsible for these schemes, but they are coordinated by the local county authorities.

In addition, there are two systems of grants for environmental measures in agriculture to encourage farmers to take steps to reduce erosion and runoff. The one system is meant to solve specific regional environmental challenges (Regional Environmental Programme, RMP) and the other system is for special measures requiring more long term investments and maintenance (SMIL).

In the SMIL system farmers can for example apply for grants to establish constructed wetlands or sedimentation ponds, hydrotechnical installations, waste water treatment facilities or re-open culverted streams. Both investment and maintenance may be granted. The local county authorities are responsible for these schemes.

In 2005, the agricultural environmental programme was changed from national to regional level. Each county has from 2005 drawn up a regional environmental programme (RMP) for the agricultural sector, reflecting the main environmental problems in the county, the local agricultural production system and the types of grants available to farmers. The county governor is responsible for the management of these schemes. Practices that may be eligible for grants include:

- Changed tillage, stubble/minimum-till rather than bare soil during the winter
- Buffer zones along streams and lakes

- Grassed water ways
- Grass on flood areas
- Catch crops
- Manure application in spring and growing season

The priority of these grants varies from county to county, and the county governors are responsible for selection of measures and the level of grants.

3.1.3 Water protection policies and management in Norway

This section is mainly based on Prestvik et al. (2013) and Øygarden et al. (2012).

3.1.3.1 Reduced tillage and other measures to reduce phosphorus losses

Norway's agri-environmental program consists of many subsidies paid to farmers who undertake certain practices or implement measures that reduce nutrient runoff to water. One part of the program is called "Runoff to water" and it was this part in particular that was evaluated in 2012 (Øygarden et al. 2012). The participation, impact and cost effectiveness was compared in 2006 and 2011 in 9 out of 19 counties in Norway. Since 2005, the agri-environmental program have been regional in nature, which means that county authorities can adjust measures to suit regional conditions like agricultural production, erosion risk and pollution level. Counties have the freedom to choose level of payments, adjust measures and implement new measures.

In 2010, 56.3 % of the area used for cereal production is associated with changed tillage. The measure "changed tillage" comprises several tillage practices where the reference practice is ploughing in the autumn after harvest. Changed tillage then refers to four practices: harrowing in autumn, direct drilling of winter crops, no till in autumn but ploughing or harrowing in spring and direct drilling in spring. These tillage measures were implemented on a larger area in 2010 than in 2006 in all but two counties. Implementation of other measures, e.g. grassed water ways, grassed buffer zones along streams and areas converted to permanent grassland, also increased from 2006 to 2010.

In Norway, cultivated area is classified according to erosion risk where category 4 has the highest risk. Measures like changed tillage and conversion to grassland have the highest potential to reduce P losses when implemented on soils with the highest risks of erosion. In one county, changed tillage in autumn or conversion to grassland is implemented on all areas with erosion category 3 and 4. In the other counties, the area in erosion category 3-4 that are ploughed in autumn varies from 20 to 57 %. From 2006 to 2010, the area of changed tillage in autumn increased the most in erosion category 1 and 2. Payments to farmers are differentiated according to erosion risk, but level of payment varies between counties. The evaluation found no evidence that the differentiation results in an increased implementation on areas with higher risk of erosion. Size of payment is only one of many factors that affect farmers' decision to implement measures.

A survey of farmers' attitudes and knowledge revealed that farmers demand counselling because of local variations when they decide what measures to implement (Refsgaard et al. 2010 and 2013). Farmers' attitudes and knowledge may be an important factor in adopting changed tillage practices. In catchments

where focus has been on changed tillage practices for several years, farmers' adoption rate of these practices is higher. This suggests that farmers' awareness of this practice affects the rate of adoption. Increased need of pesticides has been related to changed tillage practices. That and other believed or real negative effects may also stop farmers from adopting such practices.

Some catchments have additional requirements where 60 % of total cultivated area should have no-till in autumn, direct drilling of winter crops or permanent grassland. These requirements were fulfilled in 2010 which makes the rate of implementation much higher in catchments with these requirements compared to other catchments in the same county.

These catchments have also implemented more measures like buffer zones, grass-covered buffer zones along streams and flood prone areas. Estimated effect on erosion from the changed tillage measures is 290.000 tons, which means a 9 % increase in total reduced erosion. Net effect has increased in two counties because total area with no till in autumn has increased. In another county net effect has increased despite reduction in total area with changed tillage because such practices is implemented on a larger share of area with high risk of erosion. Reduction in erosion per payment also varied across counties but had a total reduction because there was an increase in payments for areas with low erosion risk.

According to the evaluation, there is still potential for reduced P leaching by reducing tillage on land with high risk of erosion. In particularly eutrophication sensitive areas, additional measures may be necessary and the development of a "packages" of measures that are suited for local conditions may be a natural extension of the agri-environmental payments.

The implementation of the EU WFD will increase the need for local effective measures. Areas with high risk of erosion and/or that are particularly sensitive to eutrophication may need locally adjusted packages that contain several measures that are supported by payments to compensate farmers for extra costs or loss of income. At present, there is little coordination between measures under different programs that can potentially reduce nutrient losses, e.g. changed tillage and fertilizer planning. In some areas, reductions in applied P and improved utilization of animal manure may be effective measures to reduce P surpluses. Reduction in applied fertilizer can also have other positive environmental effects like reduction in nitrous oxide emissions.

With climate change the need for measures to reduce nutrient leaching may be increasing, especially with increases in rainfall and higher temperatures that increases risk of erosion. The agricultural sector must adapt to climate changes and at the same time reduce GHG emissions and nutrient leaching. Agri-environmental program should include all environmental issues so that conflicting effects of measures does not arise.

3.1.3.2 Water governance

When a significant change in water quality is required, the governance of the catchment may be as important as measures and agri-environmental payments. The Lake Morsa in south-eastern Norway is an example of how local authorities can team up with stakeholders and return a lake to good ecological status, as is the aim of the WFD (Refsgaard and Gunnarsdottir 2012). The watershed included 9 municipalities in two counties and served as source of drinking water and recreational area for around 65.000 people. However, heavy loads of P lead to eutrophication and toxic algae blooms. In 1999 the Morsa river basin

organization was established and a process of creating trust and collaboration between the stakeholders were started.

The Morsa river basin team based water management on knowledge, which led to public understanding and consensus. Objectives were based on analysis carried out by neutral institutes. Every municipality developed a plan for waste water treatment by 2002 and a partly regional and partly municipal environmental program for the agricultural sector came the same year and were adopted by the municipalities in 2003.

The western part of the lake required special measures to reduce P loads. All stakeholders, including farmers were invited to participate in creating an action plan. The solution was environmental contracts between farmers and the county governor where payments were given to farmers who reduced P application on their fields and implemented other measures. 73 % of farmers signed the contract and total use of P fertilizer was reduced by 75 %. No-till practices were adopted in the autumn, buffer zones were constructed along all streams and 16 wetland sediment traps were constructed. The process changed the farmers' attitudes and engagement in improving water quality.

The result was that the lake again became suitable for swimming in 2008. The result did not come without a cost. In total EUR 90 million were spent on measures, 20 million in the agricultural sector. The area of changed tillage was increased from 30 to 80 percent of the area, P fertilization was reduced by 50 % and around 70 wetlands were constructed. However, good governance of the watershed that created trust, public understanding and collective action were critical factors for implementing the measures that resulted in significantly improved water quality in the lake and rivers.

3.1.4 Level of implementation and effect

Each year Norwegian statistics (SSB) publish time series of statistic information on farmer's behavior and choices in an annual report (Bye et al., 2015). Selected data and text from this report related to nutrient application, soil tillage and specific mitigation methods is presented here below.

3.1.4.1 Fertilisers and manure

The number of domestic animals, and thereby the quantity of manure and its nutrients, has decreased during the last ten years (Bye et al., 2015). In 2013, the number of animal manure units was calculated to 860 000. An animal manure unit (AMU) is a unit for livestock defined according to the amount of P secreted as excrement and urine (14 kg P/AMU). One calculated animal manure unit is equal to 1 dairy cow, 3 breeding pigs, 7 winter-feed sheep/goats or 80 hens etc. Measured by nutrient content at the national level, 33 per cent of all N and 58 per cent of all P used in the agriculture come from animal manure (Bye et al., 2015). These numbers, however, differ heavily between counties with the least available animal manure in the south-eastern counties.

From 1980 the sales of N have been quite stable, while the sales of P and potassium have decreased significantly, which among other may be due to high rise in prices and changes recommended nutrient application rates. In 2012/2013 the sales of commercial fertilisers were 432 000 tons, about 1 per cent

more than in 2011/2012. The sales of nitrogen were 97 000 tons and the sales of P were 8600 tons. In 2010, the sales of N in fertilizer dropped in Norway due to increased prices (Figure 3.1.1).




Figure 3.1.1. Sales of N fertiliser in the Nordic-Baltic countries 1989-2010 (FAO; Bechmann et al., 2014).

3.1.4.2 Implementation and effects of measures

The EU WFD, which Norway must apply to through the EEA-agreement, divides the country into water regions. The main purpose of the directive is to achieve “good conditions” in all waterways as regards to pollution and ecological conditions.

The county councils within a watershed are Water Region Authorities and they are responsible for the regional management plans including environmental goals and cost indications for the proposed measures for the watershed in focus (<https://www.regjeringen.no/no/aktuelt/fylkeskommunen-ny-vannregionmyndighet/id589475/>). The plans must be agreed on by the county government and approved by the Ministry of Climate and Environment. Following this the individual measures will be processed in the sector authorities and this includes a more specific evaluation of advantages and disadvantages.

Total contributions of nutrients from Norway to the North Sea are described by Selvik et al. (2015). The size of human discharges of nutrients – P and N – from agricultural activities into the waterways and oceans vary markedly between the different water regions. The water regions Glomma and Vest-Viken are the two regions where agriculture accounts for the largest relative contribution of total losses, 38 and 39 % of P losses, and 41 and 30 % for N respectively. Aquaculture is almost non-existing in these regions. Thus agriculture ranks high in relative contributions of discharges in the south-eastern areas of the country.

In 2013, the area of cereal constituted 0.30 mil. ha, or 29 % of the total agricultural area in use. Soil tillage methods in cereal areas are highly important for the risk of erosion and the risk of phosphorus losses from these areas to the water bodies. Autumn ploughing has been shown to increase both erosion and P losses (Bechmann et al., 2014). From winter wheat fields, which are ploughed before drilling, the highest P losses have been registered, but also in spring cereals autumn ploughing cause the highest P losses (Figure 3.1.2).





Figure 3.1.2 Effect of different soil tillage methods on P loss relative to autumn ploughing (Bechmann et al., 2011).

The traditional soil tillage method until 1990 was autumn ploughing. In 2000, the area of cereal ploughed in autumn covered 52% of the total cereal area. In 2013, the area with cereals ploughed in the autumn covered 46% of the total cereal area (ploughed in autumn 2012). Subsidies of NOK 164 mill. were given to change tillage methods, including catch crops and grassed water ways in 2013 (Regional environmental program) (Figure 3.1.3).

Figur 11.15. Korn- og oljevekstareal, etter metode for jordarbeiding og tidspunkt for såing¹. 2000/2001–2012/2013. Prosent




¹ Haustsådd korn er endra frå innhaustingsår til hausten det blei sådd.
Kjelde: Statens landbruksforvaltning og Jordbruksstatistikk, Statistisk sentralbyrå.

Figure 3.1.3 Trends in soil tillage methods (Bye et al., 2015; In Norwegian).

The cereal area which is sown in autumn varies a lot from year to year. If one look at the period 2000-2012, the least area was sown in autumn in 2011 with 1.2% of the cereal area, and the most was autumn-sown in 2006 with 14.4%. For 2013 the number seems to be 4.2% (Figure 3.1.3).

During the period from 1990 to 2002 Norwegian Statistics collected data on soil tillage through “Selected counting’s for agriculture”. Through this period the autumn ploughed area was reduced from 82% to 43% of the cereal area. Furthermore, it was shown for 2010 that 104 000 ha or a bit less than 34% of the cereal area was ploughed in the autumn 2009 and had no plant cover during winter. Approx. 6% or 20 000 ha of the cereal area was harrowed in autumn 2010. 86 % of the autumn harrowed area was sown with winter wheat, had catch crops or was covered with plant residues. In total 113 000 ha or 37% of the cereal area had no plant cover during winter 2009/10. Mitigation measures, such as catch crops and grassed water ways received special subsidies from autumn 1991. When the Regional environmental program started in 2005 vegetated buffers were also included in these subsidies. In 2012, subsidies were given for 424 km grassed water ways, 1232 km vegetated buffers and 5770 ha of grassed environmental area. The total subsidies for this were NOK 23.3 mill. The corresponding number for 2005 was NOK 4.1 mill.

Figur 11.16. Areal med fangvekstar. Heile landet og utvalde fylke. 2000, 2005, 2010, 2011 og 2012. Prosent



Kjelde: Statens landbruksforvaltning.

Figure 3.1.4. Trend in area with catch crops for counties and the whole country, 1 dekar = 1/10 ha (Bye et al., 2015; In Norwegian).

The area with catch crops reached its top in 2002 covering in total 35 000 ha with subsidies amounting NOK 37.7 mill (Figure 3.1.4). Later this area has gradually declined and was in 2012 4400 ha. The amount of subsidies per area decreased simultaneously. In 2012, subsidies were given to catch crop in cereal areas, early potatoes, vegetables and others. The rules for subsidies vary between counties.

Figur 11.17. Talet på nye fangdammer og våtmarker og akkumulerte tal for fangdammer og våtmarker som det er gitt tilskot til. 1994-2012




Figure 3.1.5. Trends in number of sedimentation ponds and constructed wetlands (Bye et al., 2015; In Norwegian).


In total NOK 67 000 was given to upgrading of 22 sedimentation ponds and constructed wetlands in 2012 (Figure 3.1.5). The county Buskerud had the most with 18 sites for NOK 27 000 in total. In Rogaland county 4 sites received NOK 40 000 in total. Establishment of sedimentation ponds and constructed wetlands are nature based systems to reduce runoff of soil articles and P. Subsidies for establishment of sedimentation ponds and constructed wetlands are part of the SMIL-system. This system was started in 1994 and since then the number of new sedimentation ponds increased, especially from 2000 to 2002 when the number increased from 39 to 100 per year. In 2012, NOK 3.1 mill. was given in subsidies for establishment of 38 new sedimentation ponds and constructed wetlands. During the period from 1994 to 2012 subsidies for in total 941 sedimentation ponds and constructed wetlands has been given. The number is especially high in the Rogaland County. In 2012 subsidies were given for 17 sites in Rogaland and 4 sites in Østfold and Oppland.

To reduce erosion and nutrient runoff, SMIL-subsidies are also given for hydrotechnical installations. In 2012, NOK 26.8 mill. was given to 592 hydrotechnical installations (Figure 3.1.6). The counties with the most arable land receive the most money for hydrotechnical installations. Akershus county accounted for NOK 8.9 mill. for 172 sites and county Østfold accounted for NOK 5.9 mill. for 127 sites.

In areas where cereal- and vegetable-production is dominating some of the most important measures are changed tillage, vegetated buffer zones, grassed waterways, catchment crops after harvesting, maintenance of catchment ponds and in special areas also environmental contracts. Subsidies for these measures are given through the RMP. Subsidies for investment measures such as establishment and maintenance of sedimentation ponds, and hydrotechnical installations were given through SMIL. In 2014 around 205 mill. NOK was given in subsidies to environmental measures shared between 159 mill NOK through RMP and around 46 mill. NOK through SMIL (Figure 3.1.7). The last years there has been an increase in SMIL measures and from 2010 until 2014 the total subsidies increased from 26 to 46 mill NOK.

Figure below shows the percentage distribution of subsidies within RMP and SMIL for water environmental measures.

Figur 11.18. Utbetaling og supplering av hydrotekniske anlegg i SMIL-ordninga. 2001-2012. Tilsegsbeløp og tal anlegg



Kjelde: Statens landbruksforvaltning.


Figure 3.1.6 Improvement of hydrotechnical installations (subsidies and number of sites) (Bye et al., 2015; In Norwegian).

Cost related to changes in tillage practices was estimated as the change in farmers' gross margin as a result of the change. An analysis of farmers' gross margins with different tillage practices in two counties in Norway, found that changing from winter crops to spring crops with changed tillage does reduce farmers' income (Refsgaard et al., 2010 and 2013). However, there are significant variations in these costs. In some areas, the payment covers the cost while not in all areas. The costs of reducing 1 kg of P ranged from NOK 2 000-3 000 on land with low erosion risk, to NOK 200-300 on land with great risk of erosion. Payments for changed tillage in areas with low risk of erosion are not considered cost-efficient, i.e. loss in income for farmers exceeds the environmental benefit of reduced erosion.

Tilskot til vassmiljøtiltak i 2014

Regionalt miljøtilskot

Spesielle miljøtiltak i jordbruket



* Tal for RMP i 2014 er førebelse

** Andre tiltak er vegetasjonssoner, oppsamlingsanlegg for avløp og gruppa anna

*** Gjeld òg stubb i område med jordtap og flomutsette område

Figur 21: Prosentvis fordeling av tilskot til vassmiljøtiltak for å hindra avrenning frå jordbruket i SMIL og RMP i 2014. Totalt beløp for tilskot til vassmiljøtiltak i RMP er 159 mill. kroner og for SMIL er det 46 mill. kroner.

Figure 3.1.7 Percentage of subsidies to reduce nutrient runoff from SMIL and RMP in 2014 (Trøtscher et al., 2015).

3.1.4.3 Cost-effectiveness of measures

The cost-effectiveness of mitigation methods has been included as an important criterion for selection of mitigation methods to be included in the PoM in the WFD. Refsgaard and Bechmann (2015) evaluated the cost-effectiveness of various soil tillage methods for different counties and areas in Norway. The most striking results were the large variation in cost-effectiveness related to variation in erosion risk, with the best cost-effectiveness obtained by implementing mitigation measures on high risk areas of erosion. Furthermore, Refsgaard and Bechmann (2015) found that spring harrowing was the cheapest way to reduce P loss, followed by autumn harrowing and spring ploughing in spring cereals. However, the costs of spring harrowing may not cover the long-term effect on weeds and pests by avoiding ploughing.




Figure 3.1.6. Cost-effectiveness for spring ploughed compared to autumn ploughed spring cereals for different counties in Norway (Refsgaard and Bechmann, 2015).

Implementation of changed tillage methods was more cost-effective, that is cheaper, in spring cereals compared to winter wheat. The difference in cost-effectiveness between counties was especially evident for spring harrowing and spring ploughing because of differences in soil type and related agronomic options. The cost-effectiveness of mitigation methods is an important criterion in water management plans, but when comparing the cost-effectiveness between different sectors, one also needs to consider the differences in type of P lost, variations due to the differences in soil type, agronomic conditions and socio-economic factors. The unintended negative impacts on health and environment of changed tillage and of the potential changed use of herbicides and fungicides must be considered in the management and the policy instrumentation. The study by Refsgaard and Bechmann (2015) provides considerable evidence for using local approaches to reduce such pollution from agriculture in a cost-effective way. A local and targeted approach implies that the environmental objectives may be reached at lower costs than a general regulatory approach that does not consider regional variation.

3.1.5 Norwegian summary

Most of the eutrophication problems in rivers and lakes are related to P and the measures implemented in Norway are especially focusing on reductions in P loading. The implementation of mitigation measures consist of general production grants (among others demanding 2-m buffers and nutrient management plan), legislation on manure management and subsidies for e.g. changed soil tillage, grassed buffers and sedimentation ponds administered through the Regional Environmental Program (RMP) and Special measures in agriculture (SMIL).

The agricultural environmental programme was changed from national to regional level in 2005. Each county has from 2005 drawn up a regional environmental programme (RMP) for the agricultural sector including practices like:

- Changed tillage, stubble/minimum-till rather than bare soil during the winter
- Buffer zones along streams and lakes
- Grassed water ways
- Grass on flood areas
- Catch crops
- Manure application in spring and growing season

The success-rate for implementation of mitigation measures was high, due to the economic incentives. Cost-effectiveness of several mitigation measures including its variation over the landscape has been evaluated (Refsgaard et al., 2013).

3.2 Denmark

The focus in the Danish environmental policies related to agricultural impact of water quality has been on reducing N losses (leakage at the root zone, emissions to air via stables) and to achieve better water quality and later with the implementation of WFD to achieve good ecological status in the waterbodies. This chapter describes the Danish regulation from the mid 80'es to now, implemented by the Aquatic action plans and the WFD. Both the action plans and the WFD implementation have been supported by intensive research programmes documenting the potential effect of measures and the implementation in the agricultural production, as well as by a well-established and efficient agricultural advisory service.

3.2.1 The Aquatic action plans (1987-2005)

The first Aquatic action plan was agreed on in 1987, following up on the "NPO" statement report from 1984. The objective of the Aquatic action plan was to reduce the diffuse N-leakage at the root zone by 50% and P-losses from point sources by 80%. These targets were remained in the second and third Aquatic action plans from 1998 and 2004. The P losses were mainly achieved by large investments in waste water treatment plants and in reducing other point sources, by reducing the P loads to the aquatic environment from industry and households. This regulation was effective and gave an immediate effect, but was also perceived as relatively costly (2,000 million per year for industry and sewage plants and more than 100 DKK/kg N) (Finansministeriet, 2001; Jacobsen et al., 2004). The costs has been distributed to Danish tax payers via the tax on water use and discharges. The N reduction measures where mainly targeted to agriculture, but since agricultural loads come from diffuse sources which are more difficult to regulate, with a time lag from the measure is implemented to the effect can be measured, the effect has taken much longer time. After the implementation of the three aquatic plans the target of 50% reduction has been reached (Mikkelsen et al., 2009; Dalgaard et al., 2014; Jacobsen et al., 2004).

One of the key policies implemented in both the Aquatic action plans is related to regulation of the N application. The N application is regulated by N-norms or quotas that set a limit for the N applied to the

specific crops grown. This N-norm includes both mineral fertilizer and the organic manure. The requirement regarding the utilization of N in manure, the manure N efficiency, is one of the highest in EU (Webb et al., 2010 (page 7)). The Aquatic action plan II from 1998 reduced the N-norms to 10% below the economic optimum. Due to changes in the optimal level the application rate is currently 18% under the economic optimum for 2014/15. The implication is that the shadow price of N is now exceeding the price of N, meaning that the farmers face incentives for a high utilization of N in animal manure. This is also supported by an obligatory requirement of N budgets and accounts. The average Danish farm use around 145 kg N (effective) / ha of which 80 kg N/ha comes from mineral fertilizer and the rest from animal manure. The use on mineral fertilizer is shown in figure 3.2.1 where the drop since the beginning of the 90's is clear.

Currently each Danish livestock holding must ensure a balance between agricultural land and the number of livestock units corresponding to a maximum of 170 kg ha⁻¹ yr⁻¹ of N from manure for cattle holdings and 140 kg ha⁻¹ yr⁻¹ of N for all other livestock holdings, which is stricter than the standard requirements of 170 kg N/ha from manure in the Nitrate Directive (Mikkelsen et al., 2010). This has kept the livestock density in Denmark at 1.1 Livestock Units (LU) per hectare, which is somewhat lower than the livestock density in other livestock intensive areas in EU (Grinsven et al., 2012). One Danish livestock unit is defined at a level of 100 kg N, measured at the storage level, and it is currently equal to 0.75 dairy cow or 4.3 sows with piglets up to 7 kg. Compared to most other EU-countries a large share of the total area is used for intensive agricultural production. Figure 2 shows the total N application and the increasing N-efficiency over time.




Figure 3.2.1. The purchase of N and P in mineral fertilizer from 1935 until 2012/13.

Source: Natur- og Erhvervsstyrelsen





Figure 3.2.2. Average consumption of mineral fertiliser and manure (left). N surplus, N efficiency and N leaching (right) for the agricultural area in Denmark, 1980-2012.

The total costs of the Danish aquatic action plans have been relative high, and since the mid 1990'ties a focus has been allocated to the cost-effectiveness of reaching the selected targets. These targets have been primarily focused on N-losses and to some degree also the P surplus. The overall costs of all the action plans so far (including agricultural and non-agricultural measures) are around €600 million (2005 prices). Of these the annual cost related to agricultural measures is about €340 million, and the rest is related to industry and sewage treatment plants. The estimated costs related to the different action plans are described in Table 3.2.1.

Table 3.2.1. Estimated costs of agricultural measures in different action plans to reduce N leaching from agriculture (2005 prices)

| | Reduced N-leaching from rootzone Tons N / year / N/ha/yr. (ex ante) | Ex-ante costs (mill. €/year) | Ex-post costs (mill. €/year) |
|---|---|------------------------------|------------------------------|
| Action Plan for aquatic environment I – API (1987) | 50.000 / 19 | 84 | Not calculated |
| Action Plan for a more sustainable agriculture (1991) | 40.000 / 15 | 134 | Not calculated |
| Action Plan II (1998 – 2003) | 48.000 / 18 | 92 | 70 |
| Action Plan III (2004-2012) | 300 / 1 | 30 | 48 |
| Total | 150.000 / 53 | 340 | --- |

Source : Jacobsen et al. (2004), Jacobsen (2004) and Jacobsen et al. (2009).

The payment of the costs in APII has been roughly evenly divided between the agricultural sector and public funding. There seems to be a tendency for the agricultural sector to pay for farm-related measures (changed production), whereas the state pays for land taken out of production through measures which are co-financed with the EU. In terms of farm types, the majority of costs, which relate to measures like changes in feeding, lower livestock density and increased utilization of N, are borne by animal farms, whereas other costs (lower N quota and catch crops) are more evenly divided among farms.

Table 3.2.2. Cost effectiveness for different measures in Action Plan II.

| | Reduction in N leaching | Total cost | Cost effectiveness |
|--|----------------------------|-------------------|-----------------------|
| | Tonne N/year | Million €/year | € / kg N |
| Wetlands | 800 | 0.7 ³⁾ | 0.9 |
| ESA-areas | 700 | 7.7 | 10.9 |
| Forestry | 800 | 4.7 ³⁾ | 5.9 |
| Organic farming | 3,700 | 14.0 | 3.8 |
| Changed feeding | 3,800 | 5.7 | 1.5 |
| Lower livestock density ¹⁾ | 140 | 1.5 | 10.4 |
| Catch crops (6 pct.) | 3,000 | 6.4 | 2.1 |
| Increased utilization of N in animal manure (15 pct.) ¹⁾ | 10,110 | 6.7 | 0.7 |
| Reduced N quota (10 pct.) ^{1) 4)} | 12,850 | 22.8 | 1.8 |
| Sum ²⁾ | 35,900 | 70.2 | 2.0 |

¹⁾ In the technical evaluation of Action Plan II the effect of these measures was not estimated individually, hence the figures given here are approximate estimates.

²⁾ Changes in use of area and animal production as well as other matters are not included in the table.

³⁾ Annuity based on 4 percent interest and infinite lifetime.

⁴⁾ A later recalculations have shown that the costs of N-norms have been higher than expected. (Kristensen and Jacobsen, 2014).

Source: Jacobsen (2004).

3.2.2 The WFD and the River Basin management Plans from 2011

The action plans have been followed up by the implementation of the WFD, where the objectives have been changed from a reduction target on N and P at the root zone, to environmental objectives of obtaining good ecological status, as in the rest of the EU and EØS countries (EU Commission, 2002). The Danish River Basin Management Plans (RBMP) are the implementation of the WFD in Denmark for the first period and describes both the present water quality, the distance to good ecological status and the measures implemented in the first planning period in the 23 catchments (Dalgaard et al., 2014). The quality evaluation show that only 28% of the streams, 22% of the lakes and 2% of the coastal waters have Good Ecological status and by 2021 this will be 54%, 44% and 42% with the current baseline projection (DØRS, 2015). More measures are needed to reach the targets by 2027.

The annual costs of the RBMP from 2011 have been estimated to be 56.6 million €. (Jacobsen, 2012). The cost of reducing the diffuse pollution from agriculture is 35 million € a year and the rest are costs to implement groundwater measures, to improve the physical conditions in streams and point source pollution. In 2012 the costs for the agricultural sector was estimated to 18 million € per year, which is somewhat higher than the expected 11 million € per year when the RBMP plans were made, due to higher costs related to the implementation of catch crops, change in norms and the ploughing of grass for feeding.

3.2.3 Measures and N-policies 2014 until today

When looking back over the years some key measures have already been highlighted such as the N-norms and the reduction to under optimal N-norms. These general regulations have reduced N-losses, but have also been costly. The costs have been relatively high for many farms as the N-norm is set according to crops, soil type and use of irrigation. As mentioned the combination with N-accounts and the requirements on utilization of N in animal manure has promoted high utilization of N in manure. The catch crops are another well used measure, which many farmers do not believe have the desired effect especially on clay soils. Today all farms have between 10-14% catch crops and some more if it is part of an increase in the livestock production.

With respect to the area related measures where area is taken out of production, wetlands and riparian zones have been discussed for many years in Denmark. Riparian zones were previously voluntary, but the area was very limited. Then in 2012 the 10 m (later 9m) riparian zones were made obligatory along all streams. The implementation turned out to cause a lot of protest among the farmers because of faulty maps associated with the decision, and the opposition against the measure led to that this measure will be abolished from 2016, although the measure has positive effects in relation to N, P, pesticides and nature aspects, if targeted to the riparian zones.

Wetland reconstruction is another example of a measure which has been implemented over many years based on co-operation between farmers and authorities, and between farmers. The first projects were implemented in 1990'ties and the total area is today over 12,000 ha. The implementation of the measure has shown large differences in costs and effects (Hansen et al, 2011), and has, among other factors, given rise to the increased focus on targeting the nutrient regulation to areas where the measures are most cost-effective.

3.2.4 Danish water councils - a new bottom-up approach to River Basin Management Planning

Most of the measures applied in Danish water management have until now been focused on command-and-control measures, mainly to ensure the effect as the voluntary measures have often not had the required effect (Dalgaard et al., 2014). With the introduction of the WFD a focus on participation of stakeholders has been put forward with an emphasis on an increased attention on the need to find new approaches with local, targeted measures (Nature and Agriculture commission, 2013). The first Danish RBMPs should have been implemented in 2009, but was firstly published in December 2014. There had been protests against the planned measures that led to a 5 year delay of the first RBMPs. The plans had over 5000 complaints and many of the main interest groups and the public weren't involved in the process. This has proven to be problematic for the Danish implementation of the first cycle of RBMP (Gertz et al., 2012). Because of the struggle with the first RBMPs, the government decided in preparing the second RBMP (2015-2021) to adopt a new bottom-up participation concept. In 2013, the former Danish government adopted a new act on water planning (Lov om vandplanlægning) (FT nr. 1606 26/12/2013). This new act on water planning allowed for a new bottom-up water management concept with the establishment of 23 water councils in all 23 River Basin Districts. The water councils were to advise the local municipalities on applying a PoMs for the improvement of the physical conditions in targeted Danish streams. This was mainly done to ensure a greater local involvement of stakeholders in the 2. cycle of River

Basin Management Plans. In an evaluation study of the water council work in 2014, this new approach has been tested and found successful in finding solutions at the local level (Graversgaard et al. 2015). The evaluation study found that the concept created legitimacy around creating PoMs and that the possibility of knowledge co-production was responsible for the process-success of this new kind of institutional arrangement and water management (Graversgaard et al. unpublished material). The question is, if the use of water councils in water policies in general would have created PoMs that would have been more cost-effective than the chosen measures and would they have managed to reach the goals?

The targets required to reach good ecological status has in Denmark been translated into a reduction of N which is required. There is much debate on these calculations and the required target. In the RBMPs from 2014 it was estimated that a further 9,000 tons of N should be reduced on top of the baseline development (Jacobsen, 2014). In 2014 a new catalogue of measures was prepared with the aim to provide a basis for choosing the most cost-effective measures (Eriksen et al., 2014). Focus in the catalogue is on the new measures as well as the most important ones, but even if there is a focus on targeted and local measures the estimates in the catalogue are still showing average measures of costs and effects, even though both costs and effects differs a lot between locations. At the same time work has been initiated to investigate new regulation paradigms to obtain more local and targeted regulation, and a new N retention map has been established (Højbjerg et al., 2015) to enable localisation of measures to the areas where the N retention is lowest, i.e. where the effect of the measure in the fjord or coastal area will be the highest possible. The new regulation paradigm, as well as the use of the retention maps, has not yet been decided on politically.

Farmers are currently most interested in measures like constructed wetlands (ponds, mini-wetlands), early sowing of winter wheat and other measures outside the farm area, such as mussel farming, compensating farming to remove N in fjords. Nature organizations are on the other hand suggesting that more area should be taken out of production to reduce N-losses and improve water quality as well as nature and biodiversity.

Denmark has a new government from June 2015 and in its 16 point plan they will abolish the N-norms, riparian zones and some other measures. It remains to be seen how this is implemented at the same time as the focus is on more targeted measures and implementation of the WFD requirements.

3.2.5 Conclusion and discussion - Denmark

It is evident that the N regulation of Danish agriculture has had a significant measurable statistical impact on the decreasing trend in N surplus, in N leaching from the root zone, reduction of nitrate in groundwater, and on the N load to coastal waters. The measures applied have succeeded in improved utilization of animal manure, fertilizer and crop rotation plans, improved utilization of feed-stuffs and on limitations on the total N application. Important factors is the knowledge-based approach with intensive research programmes documenting the potential effect of measures and the successful implementation of measures to the Danish farmers by a well-established and efficient agricultural advisory service. Cost-efficiency has been an important component for many years. The N norms and the N accounting system has both reduced inputs and improved utilization of N in manure. What we also see is that these measures have a limit. Documented significant and continuous lowering of protein content in Danish cereal have been the price for low N Norms over a number of years and the new governments 16 point plan, is part of regaining

a more competitive Danish agriculture and on the other hand choosing more targeted measures to retain a low loss of nutrient to surface waters. Therefore the future regulation is focused on trying to target the measures to areas where the effect in terms of reducing the N loads to coastal waters are the highest. Including this aspect means more targeted measures at selected locations and will give economic gains as oppose to not targeted measures. The regulatory setup to deal with this has not been decided yet.

3.3 Sweden

3.3.1 Policy

The Swedish Parliament in 1988 decided that a PoMs was to be initiated which would reduce the loss of nutrients from agriculture to the environment. There were three broad types of measures introduced to reduce nutrient losses from agricultural production; regulations, informational measures and measures supported by economic instruments. The decision was also specific with respect to the measures that were to be included in the program; manure storage capacity requirements, limited animal intensity, fees on mineral N and P, requirements for green cover during fall and winter and provision of advising to support plant demand fertilization. Initially the PoMs was driven by national environmental goals and included in national agricultural policy. Since agri-environmental policy was motivated by national interests the only obligation was to achieve Swedish national political commitments and satisfy national politicians and voters. However, membership in the EU and subsequent international agreements changed this obligation through a series of directives and treaties.

Swedish membership in the EU allowed for financing of agri-environmental measures through the EU Rural Development Program (RDP) but also required compliance with two EU Directives; the Nitrates Directive (ND) in 1996 and the WFD in 2000. In addition, as a signer of the agreement on program to reduce nutrient loads to the Baltic Sea (Baltic Sea Action Plan) Sweden committed itself to a set of quotas which specified national loads for each of the countries participating in the agreement. However, in spite of these latter commitments Swedish agri-environmental policy measures continued to have national goals as the factor which motivated their existence. These goals had been codified by the Swedish Parliament in 1990 as a set of 16 national environmental goals (Miljömål) which included several with respect to water quality. The most important of these for the impact of agriculture with respect to nutrient losses was the goal of “zero eutrophication”. Although there were additional goals which were included to motivate agri-environmental measures that would reduce nutrient loads from agricultural; flourishing lakes and streams, good quality groundwater, flourishing, a balanced marine environment, flourishing coastal areas and archipelagos, thriving wetlands and a rich diversity of plant and animal life. What is perhaps worth noting is that in the subsequent Swedish RDPs (for 2000-2006 and 2007-2013), in the descriptions of goals associated with payments for agri-environmental measures that aim at reducing nutrient losses, there were only references to the national environmental objectives. No specific reference was made to the role these measures would have for achieving compliance with the WFD or the BSAP.

3.3.2 Measures

In addition to the regulations which addressed manure handling, the other two types of measures either provided payments to landowners (farmers) for adopting practices which could reduce nutrient losses or

imposed a fee on the use of mineral fertilizers. The subsidized measures were included in the RDP while the fee system was national. In addition, funding has also been provided through government sponsored co-financing of individual local water management projects (LOVA program).

3.3.3 The tax on mineral fertilizers

The initial fees on mineral fertilizers were transformed into a tax on mineral N in the mid 1990's. Politically the taxes were not popular with the party which represented rural (farming) interests, the Center Party. In 2009 the tax was removed as part of an agreement which included higher taxes on the use of diesel fuels. At the time the tax was rescinded it was suggested that removal would have an insignificant impact on N losses. However, it was recognized that without the tax a lower price would result in a greater level with respect to the rate of economic optimal application and thereby higher leaching. The Board of Agriculture estimated that removing the tax would result in an extra 1500 tons of gross N losses and increase in loads to the sea of 900 tons. The so called insignificance of the effect was also questioned in a comment from researchers to the Board of Agriculture where it was pointed out that the reduction provided by the tax was almost equal to the reduction by most effective agri-environmental measure, cover crops (see Table 1). The tax was however repealed. Unfortunately, no measures to counteract this increase in leaching were implemented.

Some of the revenue from the tax had been used to support advisory services provided through a project, "Focus on Nutrients" jointly managed by the Board of Agriculture and the National Farmer's Organization (LRF). When the tax was repealed funding for this program was instead included as a measure under the RDP.

3.3.4 Agri-environmental measures supported by the Rural Development Programme (2000-2013)

In addition to support for the advisory program "Focus on Nutrients" included in the RDP starting in 2007-2013, other individual activities have been supported in both 2000 – 2006 and 2007 – 2013 which are expected to lead to reduced nutrient losses. The measures are quite detailed and included the following measures with a primary focus to reduce nutrient losses:

- Cover crops
- Spring tillage
- Riparian buffer zones
- Wetlands
- Extensive ley cultivation
- Controlled drainage (RDP starting in 2010)
- Buffer zones on erosion sensitive lands (RDP starting in 2010)
- Small wetlands as P traps (RDP starting in 2010).

3.3.5 Effect and implementation

It is somewhat difficult to consistently discuss the effect of the measures due to difficulties in measurement. The effect of some of the measures is either expressed as reduction in losses on fields at the

root zone level (cover crops, spring tillage, ley, controlled drainage, in field buffer zones and wetlands) and sometimes at the interface with the aquatic environment (riparian buffer zones, wetlands). The root zone estimates may not express accurate reduction due to transport time lags (groundwater, drainage water or surface runoff) and vary greatly due to soil and weather types. The effects of riparian buffer zones and wetlands are very site sensitive and in the case of wetlands, design sensitive. The effects are also categorized as gross (aggregated sum of the local effect of the measures) or net (includes a factor to account for retention between the local effect and discharge to the sea). The extensive and large area of inland lakes in Sweden and act as a sink for nutrients which means that location of some of the measures will have a large local effect which is important for compliance with the quality standards of the WFD but may have a small net effect on the sea and the nutrient load quotas of the BSAP.

Table 1. Reduced N and P loads as a result of measures in place and technological development.

| Measure | Reduction in N load | | Reduction in P load | |
|--|---------------------|-------------|---------------------|------------|
| | Gross | Net to sea | Gross | Net to sea |
| RDP supported measures | | | | |
| Focus on Nutrients | 650 | 390 | 10 | 7 |
| Cover crops and spring tillage | 600 | 375 | | |
| Wetlands | 450 | 300 | | |
| Controlled drainage | 30 | 22 | | |
| Riparian buffer zones | | | 12,5 | 6,5 |
| In-field wetlands | 100 | 50 | 8 | 5 |
| Regulations | | | | |
| Manure spreading Fall and Winter | 280 | 193 | | |
| Manure spreading along watercourses | 30 | 20 | 2 | 1,5 |
| Technological development | | | | |
| Techniques for adjusted N applications | 450 | 300 | | |
| Reduced tillage | 375 | 250 | | |
| Application of biogas waste | 180 | 120 | | |
| Total | 3145 | 2020 | 32,5 | 20 |

In 2008 the Swedish government gave an assignment to the Board of Agriculture to develop a plan of action for reducing the loss of nutrients to the environment for the period from 2011-2016 with a focus on outcomes to the year 2020. The final study report (SJV, 2010a) included an analysis of the effect of measures expected to be in place for this period as well as proposals for additional measures which if adopted could also be effective. Table 1 below reproduces the estimated reductions from the appendix to

this report (SJV, 2010b). In Table 1 the estimated gross reductions from all current measures are 3145 tons of N and 32,5 tons of P with more than half of this coming from measures supported by the RDP.


Area of buffer zones supported in the RDP 2003-2013



Implementation for some of the measures has varied over time, others have been relatively constant, some undersubscribed, some oversubscribed. The area in riparian buffer zones dipped significantly in 2008, a trend which continued into 2009 before surpassing previous levels in 2010. The explanation for the dip and subsequent recovery were the significantly lower payments in the RDP program starting in 2008 (1000 SEK/ha) compared to the previous period (3000 SEK/ha). Before the payment was raised in 2010. The operative goal for RDP enrollment was 5500 ha for the period from 2000-2006. This goal was exceeded for all the six years of the program, in 2006 there were 9080 ha enrolled (160% of the goal).

There are three practices which fall under the title of “reduced N leaching” cover crops, spring tillage and a combination of these two. Participation in all three of these follows a similar pattern from 2000-2013. First rising when support for these measures was introduced in 2000 to more than 200 000 ha before falling steadily for a couple of years and stabilizing at around 150 000 ha. The operative goal for the measures in the first LBP period were that 50 000 ha would either have cover crops and/or be spring tilled. While this operative goal was considerably exceeded (358%) there was still a potential for increasing the area. In a study over the potential of these activities to meet BSAP targets Johnsson et al (2009) estimated that there were 593 000 ha which could be included in the program, with 368 000 ha of this on lighter soils with a higher reduction potential although a large share of the lighter soils was already enrolled in the program in 2005 (around 50%) while only 10% of the area with heavier (clay) soils was in the program that year.

Åkerareal för minskat kväveläckage



The area in wetlands was supported by two types of financing; RDP and Local Investment Program (LIP). The operative goal was for 6 000 ha of wetlands to be maintained or created in the first and second LBP periods. Up through 2006 there were 3 868 ha which received program support (65% of the goal). More than two-thirds of the wetlands receiving support were small (less than 2 ha). There was also support provided for extensive ley production. The operative goal for the second LBP period was 200 000 ha a goal which was exceeded for 2007 – 2010 by around 30 000 ha each year. The Focus on Nutrients program over the period from the start in 2001 to 2012 reached out to around 8200 farm enterprises.

Although no direct comparison exists for the measures described above and the RBMPs, four measures with the potential for reducing P loads, currently supported in the RDP have been singled out by the three southern Swedish river basin districts in their Programs of Measure (Larsson, 2012):

- Small constructed wetlands for P retention (800 ha)
- Larger constructed wetlands (13 000 ha)
- Permanent grassed buffer strips (28 000 ha)
- Cover crops and spring cultivation (93 000 ha)

The same study reported that in the Northern Baltic Sea river basin district the total effect of the measures directed to agriculture is estimated to reduce gross P loads by 45 tons and gross N loads by 107 tons.

3.3.6 Measure effects and cost effectiveness

The effects of some of the measures (cover crops, spring plowing, riparian buffer zones) are based on the same models used for reporting on a regular basis (every six years) as part of the Helcom Pollution Load Compilations (PLC). These reports in turn serve as guidelines for goals and targets in both River Basin Management plans and British South Africa Police (BSAP). Therefore there is a consistency between the effects reported and the goals for these measures. The effects of wetlands on reducing nutrient losses have been the subject of an in-depth study performed for the Board of Agriculture (SJV Rapport 2010:21). These effects were also the basis for the half-time evaluation of the RDP covering the period from 2007 – 2010. The effect of the support for extensive ley is taken from information provided in the half-time evaluation. The effect of the Focus on Nutrient program is based on a recent independent evaluation of the program performed by the Agrifood Economic Center (reference).

Additional measures not included below because they are new and fewer studies of their effects have been performed. These include: structural liming, in field wetlands, in field buffer zones, controlled drainage, precision agriculture, two-stage ditches and lime filters.

3.3.6.1 Cover crops and spring plowing

The total area of the three measures included under this heading (cover crops, spring plowing and a combination of the two) were estimated in 2009 to lead to a total reduction of 1 340 tons N/yr. The average annual reduction was 9.8 kg N/ha for the area in the RDP support program. However, the estimated reduction varied across the areas of support from a low of 5.6 kg N/ha in Eastern Sweden (south of Stockholm) to 12.9 kg N/ha in the area with a lot of rain and erodible soils but with less areas eligible for support (Western Sweden). Cost effectiveness was calculated in the half-time report for the three measures by using the modeled estimates for the effects and the payments made to support each of the measures (as the support payment per hectare was different for each measure). The average cost for N reduction for the three measures in 2008 was 96 SEK/kg N. This represents a fall in average costs with respect to the average for the previous LBP period when the cost was estimated at 115 SEK/kg. The improvement in cost effectiveness may have been due to lower participation in the program in 2008 compared with the previous period due to an adjustment in support payments. The RDP from 2007-2010 included increased support for cover crops in combination with spring plowing (the most effective of the three measures) and lowered the payments for cover crops with fall plowing (the least effective measure).

3.3.6.2 Riparian buffer zones

In 2008 the area of riparian buffer zones supported with payments from the RDP was estimated to lead to a reduction of 6.5 tons of P. This results in a reduction of 0.93 kg P/ha with respect to the total hectares supported in the program. However, almost half of the total was due to commitments entered into in the previous period. Landowners who participate in the support commit to keep a buffer zone maintained for 5 years. In the new LBP the payments were lowered from 3 000 SEK/ha to 1 000 SEK which not only led to a sharp fall in the area signed up in the new program but also increased the cost effectiveness of the measure. In the previous program the P reduction was calculated at 3 250 SEK/kg while in the new program with the same effect but lower costs the reduction cost was only 1020 SEK/kg.

A new web based program, the FyrisSKZ, summarizes the cost efficiency of buffer zones along lakes, watercourses and ditches for all 12,864 sub-catchment areas of Sweden. FyrisSKZ is a public domain web

application (at <http://fyrisskz.slu.se>) which allows users to choose one or several sub-catchments from a GIS interface and view the estimated reduction of a buffer zone on P losses to the watercourse from surrounding fields, the opportunity and maintenance costs for buffer zones, and the potential area of buffer zones in the chosen sub-catchment. These estimates are presented for five individual buffer zone widths to allow the user to study the influence of the selected width on the reduction effect, costs and potential area. An application of the model ((Collentine et al, 2014) demonstrates how the model makes it possible for local authorities to raise cost efficiency through better targeting.

3.3.6.3 Wetlands

The estimated N retention of wetlands that received support from the LBP for the period from 2007-2010 varied between 49-80 kg N/ha of the support area. The estimated net load to the sea was reduced by 36-77 kg N/ha. The result then for the 747 hectares built with support over this period was between 37-60 tons N/yr. The area of LBP support is somewhat larger than the area of the wetland as it may also include surrounding areas which are effected by the wetland and which are therefore eligible for support. If the actual wetland surface area instead of the LBP supported area is used to calculate the effect, the effectiveness is doubled in size. P retention for the same period and same wetlands was estimated to vary between 0.96-5.05 kg P/ha resulting in a total reduction of 0.7-3.8 tons P/yr.

3.3.6.4 Extensive ley

This measure in the RDP consists of two separate objectives. One part of the measure targets forested areas of Sweden with the goal of increasing biodiversity. The second objective is to increase the area in ley in the most agriculturally intensive areas of Sweden with the goal of reducing nutrient losses. Evaluating the effect of the support payments on both objectives has proved to be difficult as the payments primarily serve to keep the land already cultivated in ley in that land use. If the payments were not present it is not apparent what the land would be used for and therefore difficult to assign an effect. If the measure results in less land lying fallow than it is expected that this can be considered positive with respect to reducing nutrient losses but has not been quantified.

3.3.6.5 Focus on nutrients

In 2008 the Board of Agriculture performed an evaluation of the effect of the Focus on Nutrients program on N losses on farms that participated in the advising program (Olofsson et al, 2008). The report concluded that farms in the area where the program had been active the longest (since 2001) that the area of cover crops in participating farms was 2-4% greater than the average for all farms in the area. According to the report authors there was a greater level of participation in the cover crop measures for Focus on Nutrient farms than for other non-participating farms. A more recent report (Agrifoods, 2015) the authors concluded that increased advising visits from the Focus on Nutrients program reduced N leaching and led to increased yields. The report also compared the cost of the program with the reductions and gains in yields. Each advising visit was estimated to reduce N farm surpluses by 3.6 kg/ha of which 2,12 kg/ha was attributed to the advice provided through the program and after taking into account leaching as a share of the reduced surplus and retention led to a reduction to the sea of 0,89 kg/ha. On the basis of other studies this reduction was valued at between 4.1 – 81.8 SEK/kg. With the average farm size within the program of 115 ha this led to a value of the reduction of between 420-8370 SEK per program advising visit. In 2013 the program cost 440 million SEK which gave a cost of 23 400 SEK for each farm visit. However, the effect on

increased yield values from the visit was estimated at almost 36 000 SEK which more than covers the cost of the visit.

3.3.7 Swedish summary

3.4 Finland

3.4.1 Introduction to Water protection policies of agriculture in Finland 1995-2015

This section describes the change in Finnish agriculture's water protection policies and measures. The policies are analyzed from the perspective of their objectives, coherence, evidence base, compliance, reference payment levels and evaluation. OECD (2012) provides an overview of these policy aspects.

3.4.2 Objectives and Coherence

The objectives of Finnish water protection policy are formed by national implementation of the EU directives and international commitments such as the Helsinki convention for the protection of the Baltic Sea. In addition there are national strategies and targets such as "Guidelines for water protection for 2015". However, the most important measures for non-point pollution control fall under Rural Development Programme and more specifically under the agri-environmental scheme of the Programme.

Prior joining to the EU, Finland had no scheme on reducing environmental impacts of agriculture. Voluntary environmental protection measures at farms were possible, but agricultural subsidies were targeted on increasing production. Even the objective of the fertilization taxation (1976-1994) and obligatory set-aside has been considered to be the financing of subsidies and cutting of over-production caused by crop price subsidies, rather than, or at least not solely, the water quality concern (Aakkula, 2006).

Objectives of the first agri-environmental subsidy scheme after Finland's EU membership were related to sustainable agriculture, with focus on water quality and biodiversity. Maybe not so obvious for the public, the agri-environmental scheme was also an instrument of income subsidy for the farmers (Aakkula 2006). Possibly related to this dualism between the goals of the scheme, the objectives were set at a highly general level and without specific targets regarding the quality of the water, for example a measurable range of concentrations in rivers or lakes was not used. Originally, implementing the agri-environmental scheme was assumed to reduce suspended sediment and thus also nutrient losses into water bodies by about 20–40% (Valpasvuo-Jaatinen et al. 1997).

At the more general level of water protection policy, governed by Ministry of Environment, a quantifiable target of halving the N and P loads to both inland waters and sea was set for 2005. While this government approved guideline contained more specific measures such as reduction of erosion from fields with technical solutions, improved nutrient management and handling of manure, the measures themselves did not have target levels, and the practical implementation of reaching the targets by policy intervention was left rather for the Ministry of Agriculture and Forestry, which was in charge of the agri-environmental

subsidy scheme. In the Rural Development programme, there were targets for measures in general terms such as reduction in erosion, but not their impacts on water quality.

A new Rural Development Programme of Finland was approved by the European Commission in 2000. Its objectives included reducing environmental load and sustaining agricultural production. These objectives, including production, were repeated for the agri-environmental scheme of 2000-2006. More specifically it was stated that in the longer run, measures in agri-environmental subsidy would cut down N and P loads by 50% from the levels of 1990s. Water Framework Directive (WFD), which came into force in 2000, is not mentioned in the Rural Development Programme or later in the interim report of the Ministry of Agriculture and Forestry concerning the implementation of the agri-environmental scheme (Wallenius & Kauranne 2003), while several pages are dedicated to economic outlook of the Finnish farmers. Within WFD, the agri-environmental scheme is considered as the main tool to control nutrient load from agricultural areas. Alas, conclusions of another interim rapport (Silvo et al 2002) by Finnish environment institute regarding the insufficient performance of measures relative to the targets seem to have made no mark in the agricultural policy domain.

Common Agricultural Policy (CAP) was reformed in 2000, resulting to several changes in the Finnish agricultural policy, but the objectives of the Finnish agri-environmental scheme remained generally unchanged even in the following programme period of 2007-2014. Similarly, the changes to national legislation for implementing WFD that were approved in 2004 were not explicitly accounted for in the revised agri-environmental scheme.

As the previous target year of 2005 for national water protection guidelines passed without reaching its objectives, a new set of targets for water protection was set for 2015 (VN 2006). The targeted reduction levels were also decreased. The cut in the level of nutrient loads to water was aimed to be one-third of the load levels of 2001-2005. Still no specific performance targets were defined for the measures of the scheme.

As expected, Finland adopted water management plans with the overall objective of good ecological status (Mäenpää & Tolonen 2011). To support implementation of these plans quantitative targets for number of specific measures were set by government (VN 2011).

CAP revision of "Agenda 2000" and the CAP "Health Check" agreed in the EU in 2008 were detaching further subsidies from production and introducing cross compliance requirements for good farming practices. From environmental perspective these changes improved the coherency of the main CAP income element with the agri-environmental part. However, many national subsidies in Finland were still allocated based on cultivated crops, conflicting with the aims to reduce nutrient loads in the agri-environmental scheme. The CAP "Health Check" also abolished the minimum requirements for fallow.

3.4.3 Reference level for payments, implementation and compliance

The EU membership introduced more stringent environmental regulation in Finland. Both Nitrate and Water Framework Directive were considerable changes to the national agricultural water protection policies. However, the cross compliance requirements for good agricultural practice concern a wider range of production conditions than the Nitrate directive, which focuses on N. WFD is not directly influencing the

reference level at all. Thus, the reference level for which the agri-environmental measures are to be compensated is rather dependent on CAP instead of the other policies.

Due to a high participation rate in the agri-environmental subsidy scheme the effect of cross compliance is not as much in directly changing the farmers' practices, as changing the agri-environmental subsidy scheme. For example non-cultivated strips at field edges were part of the good farming practice, and due to the cross compliance they became part of the reference level for which farmers would not be entitled to compensation in the agri-environmental scheme. The revised subsidy is calculated based on the area which exceeds the good farming practice requirement given by width of 0.6 meters in contrast to 1-3 meters required in the agri-environmental scheme depending on the type of stream, ditch or lake the field is located next to.

Similarly, for N the reference level for which the compensation depends on is set by the Nitrate directive, which is applied in the whole of Finland. The reference level itself is public, but the details of the levels of compensation are taken within the Ministry of Agriculture and Forestry. These reference level calculations have not been published, but they have been subject to an approval by the commission of the European Union and nationally influenced by various stakeholder groups that participate in the committees drafting the agri-environmental scheme.

The Finnish agri-environmental scheme was built on three types of subsidies that were both regionally and functionally differentiated. The first element was the largest one in subsidy volume and a precondition for farmers to access the other two elements that were containing most of the measures of the scheme. The first element, labelled "basic subsidy", contained maximum allowed fertilisation levels differentiated between crop and soil types as well as general compliance with good farming practice, which was later adopted from the scheme to the CAP cross compliance requirements. The basic subsidy also included information dissemination in the form of farmer training and monitoring. The second element, entitled "additional measures", was containing options from which the farmers would have been required to choose between two to four measures. The third element was called "specific subsidy" and contained measures that were completely optional and also harder to implement in the subsidy per year per hectare (of farm size) -frame, which was applied for the other two elements.

Additional measures and the fertiliser use limits were tweaked between the agri-environmental schemes, but the basic composition remained intact until the revised scheme of the latest Rural Development Programme (2014-2017). Additional measures included further adjustments to fertilisation, increasing vegetation cover of farm land during winter, cultivation of catch crops, diversification of farming, extensive grass cultivation and manure application during the growing season. The third element contained establishing of vegetated buffer zones and wetlands, as well as, ground water area specific cultivation, treatment of runoff water, organic production and production of indigenous species or traditional biotopes.

More significant than the change in the measures was the change from different farm size classes in the first programme to only one size class in the following programmes, which implicitly established an income subsidy within the agri-environmental scheme. As agri-environmental subsidies are calculated based on an average farm size and contain several fixed cost elements that do not scale up with the farm size, the bigger farms profit (and smaller farms lose out) even though the subsidy as a whole would be income neutral. If the subsidy level is set so that even the smallest farm is fully compensated, the rest of the farms

gain an income subsidy. Having several size classes meant that bigger farms were not overcompensated as much as with a scheme having a single size class.

Throughout the existence of the scheme, the participation rates have been high, nearly 90% of the farms. However, the farmer compliance has not been monitored in any of the impact assessment and monitoring reports of the scheme (Palva et al 2001, Turtola & Lemola, 2008, Aakkula & Leppänen 2014). The agri-environmental scheme was enforced as a part of the overall CAP compliance checks in which 5% of the farms were inspected. There is no public record of compliance. The committee set by the Ministry of Agriculture and Forestry considered added efforts in monitoring and enforcement as non-recommended extra burden for farmers (Wallenius & Kauranne 2003).

3.4.4 Evidence base and Policy evaluation

Agri-environmental schemes are evaluated and renewed approximately every fifth year. Due to difficulties in monitoring non-point source pollution such as nutrient runoff and leaching from agriculture, there is considerable uncertainty regarding the effectiveness of the agri-environmental measures in Finland. The measures with most expected impact on nutrient loads are fertilisation limits and erosion control by various types of tillage and vegetation cover. The effectiveness of both types of these measures has been established in empirical studies, but the overall effects on nutrient concentration in water bodies can be only measured given environmental and production changes not related to the measures themselves at a catchment scale, or modeled based on the physical processes.

Agricultural nutrient loading in Finland is evaluated by using long-term monitoring data of discharge and nutrient concentrations of 20 river basins and seven small catchments. Further, in two of the sub-basins of the large river basins farmers were interviewed annually of their cultivation practices. Altogether these river basins covered 30% of the total field area in Finland. Monitoring of catchments is based on mixed methods including continuous automatic monitoring, periodic water sampling and more detailed less frequent sample analysis.

The largest portion of the agricultural nutrient load (around 80% of the total load) was in the drainage area of the Archipelago Sea. As an average of all river basins, 50% of the total TN load originated from fields if field percentage exceeded 15%. In the case of total phosphorus (TP) the share was even higher.




Figure 3.4.1. Annual and flow normalized nutrient fluxes to the Baltic sea from the 20 river basins

Specific agricultural TN loading value was $1340 \text{ kg km}^{-2} \text{ yr}^{-1}$ and TP loading value $83 \text{ kg km}^{-2} \text{ yr}^{-1}$. The TN load from agriculture increased until the period 2000–2006, but then turned to decrease. In Finland large amount of new field has been cleared due to demand of field area in expanding dairy farms since 2000

(Luke 2013, Niskanen and Lehtonen 2014). Almost 30% of this new field area was cleared on organic soil types (Niskanen and Lehtonen 2014), which were naturally rich in N. Clearing of new field explained 50 % of the increase in the TN load to the Baltic Sea between the periods 1995–1999 and 2000–2006 (Rankinen et al. 2016). Increased trend in TN concentrations was partly explained by increased air temperature in early autumn, as higher temperature is assumed to accelerate N mineralization (Aakkula and Leppänen 2014). Partly, higher concentrations could be explained by cultivation of more N intensive crops (Lankoski and Ollikainen, 2011).

An assessment on N tax in Finland pointed out that fertilisation limits did not reduce the N fertilisation quantities for most crops compared to the average economically optimal levels in any of the Finnish agri-environmental schemes (Hilden et al 2007). However, in 1990, fertilizer use was 112 kg N ha⁻¹ and in 2007–2009, the average N use was 73 kg ha⁻¹ (Aakkula and Leppänen 2014). It seems that pre-EU national yield subsidies had a large impact on the optimal N use.

The TP load peaked already in the first program period (1995-1999) and decreased steadily since then. The decrease was 20% which reached the original target of agri-environmental scheme but was still not enough for achieving the current official targets of the load reduction to the surface waters. As climate or large scale land use changes did not explain the decrease, it can be assumed to be mainly due to success in agri-environmental scheme (Ekholm et al. 2015, Rankinen et al. 2016). For P, the decrease in load can be attributed to both reduced use of fertilisers and increased vegetation cover (period of minimum requirement for fallow, increasing trend in reduced or no-till cultivation, buffer zones and strips) as well as changes in various manure management practices (maximum animal density requirement, limited dispersal periods, trend in hose dispersal). These trends are not only following from agri-environmental or water protection policies, but influenced by non-related increases in input prices, low prices of crops and the general trend of decreasing farm numbers and increasing share of farms managed as secondary source of income.

Studies have shown that for an extensive period before the agri-environmental scheme, farmers were using higher quantities of P than what was removed by crops. This practice was supported by research indicating that the stock of P in the soil would improve the crop yield potential (Saarela 1997). More recent research has pointed out that lesser soil P stock could be economically optimal for farmers and in general the current research paradigm advocates less use of P (Ylivainio et al. 2014). Furthermore, increasing fertilisation prices might have made Finnish farmers more conscious of the crop needs. For example in 2008 global P prices peaked with 800% increase (FERTSTAT). Irrespective of the ultimate causes, the average P fertiliser quantities decreased from 30.7 kg P ha⁻¹ in 1990 to 7 kg ha⁻¹ in 2007-2009 (Aakkula and Leppänen 2014). For 2008, a drop in global fertilization sale was observed (FERTSTAT). Thus, it seems reasonable to infer that the increasing trend in fertilizer prices in Finland too, has been reflected in applied amounts.

Developments of several agri-environmental measures have shown a positive trend since 1995, and especially nutrient balances and use of commercial fertilizers decreased considerably in Finland in 1990–2006 and that of P only 7 kg ha⁻¹ (Aakkula and Leppänen 2014) for the whole country. The proportion of fallows varied between 6% and 13%. At the same time, the proportion of reduced tillage area increased from 30% to over 80% of the field area. Observed trends in nutrient concentrations or loads are often compared to those in the early 1990's when the share of fallow or set aside land was high. Even though the reason was other than water protection, the effect was positive on water quality.

The link from a lower P balance to a decreased P load is not without problems either. Negative balance will slowly deplete the soil P stock and reduce the load with delay (Ekholm 2003). High stocks take long to adjust relative to the periods in Rural Development Programmes.

The control of erosion is a much faster measure to reduce P loads. A reduction in P load could be the outcome of erosion control measures instead of slowly decreasing soil P stock. However, effectiveness of erosion control is not uniform and its performance as a P reduction measure in Finland is debated. There are several studies indicating that while reductions in P ending up to water with eroded particles do happen, simultaneously P is enriched to the surface soil and susceptible to leaching in a more biologically available dissolved form (SOURCES). The total decrease in the load thus available for algae, and particularly blue-green algae, which can be more harmful to environment, is not reduced as much as would be expected by just analyzing effectiveness of erosion control as a P abatement method.

As expected from different reduction levels of measures even in field experiments, also the watershed and nation level results depend on the sources and tools used to derive them (Puustinen, 2010, Helin & Tattari 2012). In Finland it seems that the reduction achieved by erosion control measures is overestimated. In watershed models failures to represent the share of dissolved P from the total P load correctly have led to calibrating the particle P share too high. The most recent assessment and monitoring study of the agri-environmental scheme found no statistical significance between increased vegetation cover and the decreased P loads in a river basin scale (Aakkula and Leppänen 2014). In a sub-basin scale with more detailed data, erosion control methods included in to agri-environmental scheme seemed to work in the erosion sensitive areas but not in the less sensitive areas where increase in dissolved reactive P may exceed the benefits of reduced particulate P load (Rankinen et al. 2016). Therefore focus should be on dissolved nutrients as they cause eutrophication in receiving waters. Thus, the conclusions of limited abatement potential and high abatement costs (Helin, 2013) hold, while the actual potential will be even smaller and costs higher.

The measures included into agri-environmental scheme are mainly effective in reducing nutrient loading. Further, the regulation has been tightening since 1995, e.g. manure spreading on fields in autumn has been gradually decreased though it is still at some extent allowed. Among the measures that are so far marginally applied are catch crops, which have high potential to reduce N leaching (Valkama et al. 2015). Measures outside the agri-environmental scheme include e.g. gypsum amendment (as well as structural liming) to fields, which has high potential to reduce both particulate and soluble P losses (Ekholm et al. 2012). Short term solutions should be better targeted to give the best possible benefit. In addition to these short term solutions long-term policy should concentrate on reducing areal hot-spots due to unbalanced production and options to produce environmental benefits as side effects of production.

3.4.5 Summary - Finland

Overall objectives have been set for water quality in Finland, but these objectives have not been specific regarding the implementation of water protection measures. For evaluation of the Finnish agri-environmental policy, it is important define against which objectives of the policy are evaluated, since farm income did not cease to be part of the debate concerning the agri-environmental scheme. For policies with multiple objectives, the coherency of all aspects of policy is difficult to achieve.

The most recent scheme has moved further towards environmental and water quality concerns. Compared with previous schemes, the objectives are defined more clearly in terms of specific concern over improving the state of the archipelago sea. However, the connection to actual implementation of WFD objectives is still stated on a general level, instead of well specified connections between measures and specific quality objectives. To some degree this reflects the administrative boundaries between the Ministries of Environment and Agriculture; agri-environmental subsidy scheme is governed in the Ministry of Agriculture and Forestry, while the implementation of the WFS is done under the Ministry of Environment. Partly, the more recent policies might have been designed more cautiously since past objectives have not been reached and the considerably uncertainties regarding the effectiveness of the measures.

The new policy scheme in Finland is based on the revision of CAP. In Finland the “greening” of CAP, is implemented by reducing monoculture cropping, which from the water protection perspective is an ambiguous method. Furthermore, directing subsidies to crops like rye in the new scheme is at odds with the water protection goals, since the subsidies tied to production increase the opportunity costs of water protection measures. Thus, splitting the CAP subsidies between production and environment is not a coherent policy from the water protection perspective. The agri-environmental scheme, as during the previous Rural Development Programmes, is a minor subsidy compared to the financial volume of the whole programme. The general choices in the income element of CAP thus could play larger role in the overall environmental effect of the programme and conflict with the water policy such as the nitrate or water framework directive.

4 CONCLUSION

UDKAST

5 LITERATURE

- Aakkula, J. and Leppänen, J. (eds) 2014. Maatalouden ympäristötuen vaikuttavuuden seuranta-tutkimus (MYTVAS 3) – Loppuraportti
http://www.mmm.fi/attachments/maaseutu/tutkimus/CxFzHox2r/MMM_myivas_loppuraportti_WEB.pdfBlicher-Mathiesen, G., Andersen, H.E., Carstensen, J., Børgesen, C.B., Hasler, B. & Windolf, J. (2014). Mapping of nitrogen risk areas. *Agriculture, Ecosystems & Environment* 195, 149-160.
- Blicher-Mathiesen, G., Andersen, H.E., Carstensen, J., Børgesen, C.B., Hasler, B. & Windolf, J. (2014). Mapping of nitrogen risk areas. *Agriculture, Ecosystems & Environment* 195, 149-160.
- Blicher-Mathiesen, G; Jacobsen, B.H.; Hansen, B. and Thorling, L. (2015) Danish policy measures to reduce diffuse nitrogen emissions from agriculture to the aquatic environment. Paper for OECD report on Nutrient Efficiency.
- Bye, A.S., Aarstad, P.A., Løvberget, A.I., Høie, H. 2015. Jordbruk og miljø, tilstand og utvikling 2014. Statistics Norway Report 2015/18. 145p.
- Dalgaard, T., B. Hansen, B. Hasler, O. Hertel, N. J Hutchings, B.H. Jacobsen, L. S. B. Kronvang, J.E Olesen, J.K Schjørring, I.S Kristensen, M. Graversgaard, M. Termansen, H.Vejre (2014). Policies for Agricultural nitrogen management - trends, challenges and prospects for improved efficiency in Denmark. *Jour. Environmental Research Letters. Environ. Res. Lett.* 9 (2014) 115002 (16pp) doi:10.1088/1748-9326/9/11/115002
- DØRS (2015). Økonomi og Miljø 2015. [Economy and the Environment 2015]. The Economic council. 2015. 375 p.
- Ekhholm, P., Rankinen, K., Rita, H., Räike, A., Sjöblom, H., Raateland, A., Vesikko, L., Cano Bernal, J.E., Taskinen, A., 2015. Phosphorus and nitrogen fluxes carried by 21 Finnish agricultural rivers in 1985-2006. *Environmental Monitoring and Assessment*, 187-216 DOI 10.1007/s10661-10015-14417-10666.
- Ekhholm, P., Valkama, P., Jaakkola, E., Kiirikki, M., Lahti, K., Pietola, L., 2012. Gypsum amendment of soils reduces phosphorus losses in an agricultural catchment. *Agricultural and Food Science* 21, 279-291.
- Eriksen, J., Jensen, PN, Jacobsen, BH, (editors), Means for the realization of 2nd generation water plans and targeted area regulation. Report prepared by the DCE, DCA and IFRO. DCA report 52 December 2014. Aarhus University.
- EUROOPAN PARLAMENTIN JA NEUVOSTON DIREKTIIVI 2000/60/EY, annettu 23 lokakuuta 2000, yhteisön vesipolitiikan puitteista
- FERTSTAT. Fertilisation database. FAO.
- Gertz, F., Knudsen, L., og Wiborg, I.A., 2012: Problematisk dansk implementering. *Vand og Jord*, nr. 3, 100-103, 2012.
- Graversgaard, M., Gertz, F., Wiborg, I.A., Kjeldsen, C., & Dalgaard, T. (2015): Vandråd – et nybrud i dansk vandforvaltning. *Vand og Jord*. Vol. 22. Nr. 1.
- Grinsven, H.J.M. ten Berge, T. Dalgaard, B. Fraters, P. Durand, A. Hart, G. Hofman, B.H. Jacobsen, S.T.J. Lalor, J.P. Lesschen, B. Osterburg, K.G. Richards, A.-K. Techen, F. Vertès, J. Webb, W.J. Willems (2012). Management, regulation and environmental impacts of nitrogen fertilization in northwest Europe under Nitrates Directive, a benchmark study. *Biogeosciences*, 9, 5143-5160.
- Grinsven, H.J.M., Holland, M., Jacobsen, B.H., Klimont, Z., Sutton, M.A. and Willems, W. J. (2013). Costs and benefits of nitrogen for Europe and implications for mitigation. *Env. Science and Technology*. Vol. 47, pp. 3571-3579.
- Refsgaard, K og H Gunnarsdottir 2012. Bridging the gap – holistic and local governance. *World Water Forum*, 12-16 March, Marseilles. Oral presentations.
- Hasler, B.; Hansen, L.B., Andersen, H.E: and Konrad, M. (2015). Modelling af omkostningseffektive reduktioner af kvælstoftilførsler til Limfjorden. Notat. DCE. Århus Universitet.
- Hansen, LB, Källström, MN, Jørgensen, SL & Hasler, B 2011, Vådområders omkostningseffektivitet: En erfaringsopsamling og analyse af omkostningerne ved at gennemføre vådområdeprojekter under

- vandmiljøplanerne VMPI og VMPII. Danmarks Miljøundersøgelser, Aarhus Universitet. Faglig rapport fra DMU, nr. 835
- Højberg A.L, Windolf J., Børgesen C.D, Troldborg L., Tornbjerg H, Blicher-Mathiesen G., Kronvang B., Thodsen H. & Ernsten V. 2015: National kvælstofmodel. Oplandsmodel til belastning og virkemidler
- Helin, J. 2013. Cost-efficient nutrient load reduction in agriculture - short-run perspective on reducing nitrogen and phosphorus in Finland. PhD Dissertation. University of Helsinki
- Helin, J. and Tattari, S. 2012. How much can be gained by optimizing nutrient abatement spatially -Cost-efficiency comparison of non-point arable loads from different Finnish watersheds. Food Economics.
- Hilden, M. Anni Huhtala, Kauko Koikkalainen, Maria Ojanen, Juha Grönroos, Janne Helin, Mika Isolahti, Minna Kaljonen, Arjo Kangas, Hannu Känkänen, Markku Puustinen, Tapio Salo, Eila Turtola, Risto Uusitalo. 2007. Verotukseen perustuva ohjauk maatalouden ravinnepestöjen rajoittamisessa. YMPÄRISTÖMINISTERIÖN RAPORTTEJA 15
- Horisontaalinen maaseudun kehittämissuunnitelma. Manner-Suomea koskevat EU:n yhteisen maatalouspolitiikan liittämissuunnitelman vuosille 2000 – 2006
<http://www.mmm.fi/attachments/maaseutu/maaseudunkehittämissuunnitelmat/horisontaalinenmaaseudunkehittämissuunnitelma/5IK0r8BYB/horisontaaliohjelma.pdf> Jacobsen, B.H. (2004). Vandmiljøplan II – økonomisk slutevaluering. [Economic evaluation of Action Plan on the Aquatic Environment II]. Report no. 169. Danish Research Institute of Food Economics.
- Jacobsen, B.H. (2004). Vandmiljøplan II – økonomisk slutevaluering. [Economic evaluation of Action Plan on the Aquatic Environment II]. Report no. 169. Danish Research Institute of Food Economics.
- Jacobsen, B.H.; Hasler, B. and Hansen, L.B. (2009). Økonomisk midtvejsevaluering af Vandmiljøplan III [Economic midterm evaluation of Action Plan on the Aquatic Environment III]. Note. Institute of Food and Resource Economics, University of Copenhagen and National Environmental Research Institute, University of Aarhus.
- Jacobsen, B.H. (2012). Analyse af landbrugets omkostninger ved implementering af vandplanerne fra 2011. [Economic evaluation of the Agricultural costs related to the River Basin Management Plans from 2011]. Note. Danish Research Institute of Food Economics, University of Copenhagen.
- Jacobsen, B. H. (2012). Analyse af omkostningerne ved en yderligere reduktion af N-tabet fra landbruget med 10.000 tons N. Notat til N-udvalget. Fødevarøkonomisk Institut, KU. FOI Udredning 2012/26
- Jacobsen, B.H. ; Abildtrup, J.; Andersen, M., Christensen, T.; Hasler, B.; Hussain, Z.B.; Huusom, H.; Jensen, J.D.; Schou, J.S. og Ørum, J.E. (2004). Omkostninger ved reduktion af landbrugets næringsstoffab til vandmiljøet – Forarbejde til vandmiljøplan III. Rapport nr. 167. Fødevarøkonomisk Institut.
- Jacobsen, B.H. (2009). Farm economic consequences of The Water Framework Directive in Europe. Journal of Farm Management. Vol. 13, No. 10, pp. 687-696.
- Jacobsen, B. H. og Steen-Kristensen, E. (2013). Landbrugets omkostninger ved den nuværende normreduktion. Notat fra Københavns og Århus Universitet. Udredningsnotat fra IFRO, KU. Nr. 14/2013.
- Jensen, C.L.; Dubgaard, A.; Jacobsen, B.H.; Olsen, S.B. and Hasler, B. (2013). A practical CBA based screening procedure for identification of river basins where the costs of fulfilling the WFD requirements may disproportionate – applied to the case of Denmark. Journal of Environmental Economics and Policy. Vol. 2, No. 2, 164–200.
- Jensen, P.N., Jacobsen, B.H.; Hasler, B. Rubæk, G. og Waagepetersen, J. (2009). Notat vedr. virkemidler og omkostninger til implementering af vandrammedirektivet.. Rapport udarbejdet til Virkemiddeludvalg II for By- og Landskabsstyrelsen. Lankoski, J. and Ollikainen, M. Counterfactual approach for assessing

- agri-environmental policy: The case of the Finnish water protection policy
<http://ageconsearch.umn.edu/bitstream/98993/2/lankoskiolikainen.pdf>
- Luke, 2013. Agricultural statistics. The Natural Resources Institute Finland,
<http://statdb.luke.fi/PXWeb/pxweb/en/LUKE/>
- Mikkelsen, S.; Iversen, T.M; Jacobsen, B.H. and Kjær, S.S. (2010). Danmark – Europe. Reducing nutrient losses from intensive livestock operations. In Gerber, P; Mooney, H. and Dijkman (2010). Livestock in a changing Landscape. Volume 2: Experiences and regional perspectives. Chapter 8, pp. 140- 153. Island Press. ISBN: 9781597266734, 208 p
- Miljøstyrelsen 1998: Dansk miljøeksport af produkter og rådgivningsydelser til vandsektoren. Miljøministeriet. . Webb J., Sørensen P., Velthof G., Amon B., Pinto M Rodhe L, Salomon E, Hutchings N., Burczyk P, J. Reid. 2010: Study on variation of manure N efficiency throughout Europe. EU Commission – Directorate General Environment. ENV.B.1/ETU/2009/0026
- Mäenpää, Milla and Tolonen, Sini. (eds) 2011. Kooste vesienhoitoalueiden vesienhoitosuunnitelmista vuoteen 2015. Suomen ympäristö 23/2011
- Niskanen, O., Lehtonen, E., 2014. Maatilojen tilusrakenne ja pellonraivaus Suomessa 2000-luvulla. MTT Raportti 150, p. 28.
https://helda.helsinki.fi/bitstream/handle/10138/37044/SY_23_2011.pdf?sequence=3
- OECD 2012. Evaluation of Agri-environmental Policies Selected Methodological Issues and Case Studies. OECD. DOI:10.1787/9789264179332-en
- Pietiläinen, O.-P., Räike, A., 1999. Typpi ja fosfori Suomen sisävesien minimiravinteena. Suomen ympäristö, ympäristönsuojelu 313., p. 64.
- Prestvik, A. S., Kvakkestad, V. and Skutevik, Ø. 2013. Agriculture and environment in the Nordic countries. Policies for sustainability and green growth. TemaNord 2013:558, Nordic Council of Ministers.
- Puustinen, M., Turtola, E., Kukkonen, M., Koskiahio, J., Linjama, J., Niinioja, R. and Tattari, S. 2010. VIHMA— A tool for allocation of measures to control erosion and nutrient loading from Finnish agricultural catchments. Agriculture, Ecosystems & Environment, 138:3–4, 15 p. 306-317
- Räike, A., Pietiläinen, O.-P., Rekolainen, S., Kauppila, P., Pitkänen, H., Niemi, J., Raateland, A., Vuorenmaa, J., 2003. Trends of phosphorus, nitrogen and chlorophyll a concentrations in Finnish rivers and lakes in 1975-2000. the Science of the Total Environment 310, 47-59.
- Rankinen, K., Keinänen, H., Cano Bernal, J. E. 2016. Influence of climate and land use changes on nutrient fluxes from Finnish rivers to the Baltic Sea. Agriculture, Ecosystems and Environment. 216:100-115. In Print.
- Rankinen, K., Gao, G., Granlund, K., Grönroos, J., Vesikko, L., 2015b. Comparison of impacts of human activities and climate change on water quantity and quality in Finnish agricultural catchments. Landscape Ecology DOI 10.1007/s10980-014-0149-1. Rural Development Programme for Mainland Finland 2007-2013.
http://www.mmm.fi/en/index/frontpage/rural_development/Rural_development_programmes/strategy_mainland_2007_2013.html
- Ylivainio K. 2010. Sarvi, M., Lemola, R., Uusitalo, R. and Turtola, E. 2010. Regional P stocks in soil and in animal manure as compared to P requirement of plants in Finland.
- Refsgaard, K. and Bechmann, M. 2015. Cost-effectiveness of tillage methods to reduce phosphorus loss from agricultural land. Journal of Environmental Planning and Management (In press).
- Refsgaard, K, M. Bechmann, AG. B. Blankenberg, S. Skøien, A. Veidal. (2010): «Kostnadseffektivitet for tiltak mot fosfortap fra jordbruksarealer i Østfold og Akershus». NILFrapport 2010-2.
- Refsgaard, K. M. Bechmann, AG. B. Blankenberg, V. Kvakkestad, A. Ø. Kristoffersen and A. Veidal (2013): Evaluering av tiltak mot fosfortap fra jordbruksarealer i norge. NILF-rapport 2013-3.
- Silvo K. M. -L Hämmäläinen and K. Forsius and T. Jouttijärvi and T. Lapinlampi and E. Santala and E. Kaukoranta and S. Rekolainen and K. Granlund and P. Ekholm and A. Räike and K. Kenttämies and A. Nikander and J. Grönroos and E. Rönkä. 2002. Päästöt vesiin 1990-2000. Vesiensuojelun tavoitteiden

https://helda.helsinki.fi/bitstream/handle/10138/40848/SYKEmo_242.pdf?sequence=1

- Tamminen, T., Andersen, T., 2007. Seasonal phytoplankton nutrient limitation patterns as revealed by bioassays over Baltic Sea gradients of salinity and eutrophication. *Marine Ecology Progress Series* 340, 121-138.
- Trötscher, T.A. and Hundere, G.C.T. 2015. Miljøstatus i landbruket for 2014. Tematisk gjennomgang av miljøstatus og virkemiddelbruk. Landbruksdirektoratet rapport 9/2015.60p.
- Valpasvuo-Jaatinen, P., Rekolainen, S., Latostenmaa, H., 1997. Finnish agriculture and its sustainability: environmental impacts. *Ambio* 26, 448-455.
- Valtioneuvosto. Vesiensuojelun tavoitteet vuoteen 2005. Valtioneuvoston periaatepäätös
- Valtioneuvosto. Vesiensuojelun tavoitteet vuoteen 2015. Valtioneuvoston periaatepäätös
- Valtioneuvosto 2011 Vesienhoidon toteutusohjelma 2010–2015. Valtioneuvoston periaatepäätös
- Wallenius S., and Kauranne, L-M. 2003. Maatalouden ympäristötuen seurantaryhman väliraportti. http://www.mmm.fi/attachments/mmm/julkaisut/tyoryhmamuistiot/2003/3v0FcYCSR/trm_2003_7_Maatalouden_ymparistotuen_seurantaryhman_valiraportti.pdf
- Väisänen, S. and Puustinen, M. (eds). 2010. Maatalouden vesistökuormituksen hallinta- Seuranta, mallit ja kustannustehokkaat toimenpiteet vesienhoidon toimenpideohjelmassa. Suomen ympäristö 23 https://helda.helsinki.fi/bitstream/handle/10138/37988/SY_23_2010_s_1_26.pdf?sequence=1
- Valkama, E., Lemola, R., Känkänen, H., Turtola, E., 2015. Meta-analysis of the effects of undersown catch crops on nitrogen leaching loss and grain yields in the Nordic countries. *Agriculture, Ecosystems & Environment* 203, 93-101.
- Ylivanio K. 2010. Sarvi, M., Lemola, R., Uusitalo, R. and Turtola, E. 2010. Regional P stocks in soil and in animal manure as compared to P requirement of plants in Finland.
- Øygarden, L., Grønlund, A., Skøien, S., Refsgaard, K., Krokann, K., Nordskog, K., Bechmann, M. 2012. Evaluering av Regionale Miljøprogram (RMP) 2011. Bioforsk rapport 7/21.

6 DANISH APPENDIX : SUMMARY OF CATALOGUE OF MEASURES (ERIKSEN ET AL., 2014)

I Tabel 0.1 er samlet en oversigt i meget kort form om de enkelte virkemidler. Desuden er der gengivet nogle af de vigtigste forklaringer og forbehold, som hører med i vurderingen af de enkelte virkemidler. Det er imidlertid nødvendigt at læse den fulde beskrivelse af virkemidlerne for at få et samlet billede af virkning, potentiale, forbehold osv.

| Virkemiddel | Referencepraksis/ kommentar | Årlig effekt 1) | N- Sikkerhed ift. N-effekt | Budget-økonomisk omkostning kr./kg N 2) | Velfærds-økonomisk omkostning kr./kg N 2) |
|--|---|--|----------------------------|---|---|
| Efterafgrøder | Jord uden efterafgrøde | 12-45 kg N ha ⁻¹ 3) | *** | 5 - 19 157 - 236 4) | 6 - 25 209 - 311 4) |
| Mellemafgrøder | Vintersæd uden mellemafgrøder | 9-13 kg N ha ⁻¹ | ** | 30 - 36 | 39 - 48 |
| Afgrøder med høj N-optagelse: > Sukkerroer > Græs og frøgræs > Foderroer | Jord uden efterafgrøde Jord uden efterafgrøde (Mangler data) | 12-45 kg N ha ⁻¹ >12-45 kg N ha ⁻¹ - | ** (samlet bedømmelse) | -114 - (-156) IV IV | -152 - (-206) IV IV |
| Tidlig såning af vinterhvede (7. september) | Normal såning af vinterhvede (23. september) | 5-8 kg N ha ⁻¹ | ** | -80 - 54 | -106 - 72 |
| Flerårige energiafgrøder | Kornrige sædskifter under den nuværende regulering | 34-51 kg N ha ⁻¹ 3) | *** | -45 - 107 | -60-142 |
| Brak (ikke permanent udtagning) | Jord i omdrift | 35-58 kg N ha ⁻¹ 3) | * | 28 - 190 | 37 - 253 |
| Permanent udtagning | Jord i omdrift | 50 kg N ha ⁻¹ | ** | 69-83 | 91-130 5) |
| Randzoner | Jord i omdrift og varig græs | 37-74 kg N ha ⁻¹ | * | 47-93 | 62-123 7) |
| Fjernelse af biomasse i randzoner og engarealer | Ingen fjernelse af biomasse | Pt. ikke datagrundlag | IV | IV | IV |
| Skovrejsning | Jord i omdrift | 50 kg N ha ⁻¹ | ** | 50-153 6) | 66-203 6) |
| Forbud mod | Jord der | 10 kg N ha ⁻¹ | ** | 1 | 1-2 |

| | | | | | |
|-----------------------------------|------------|--|--|--|--|
| jordbearbejdning i visse perioder | bearbejdes | | | | |
|-----------------------------------|------------|--|--|--|--|

Årlige kvælstofeffekter i form af estimeret, reduceret kvælstofudvaskning, sikkerhed i forhold til estimeret kvælstofeffekt, samt budget- og velfærdsøkonomiske omkostninger for hvert virkemiddel. IV og IR angiver, at værdien er henholdsvis ikke vurderet eller ikke relevant.

| | | | | | |
|---|--|---|----|----------|----------|
| Forbud mod omlægning af fodergræs om efteråret | Intet forbud mod omlægning om efteråret | 36 kg N ha ⁻¹ | * | 14 | 18 |
| Reduceret jordbearbejdning | Konventionel jordbearbejdning | 0 8) | ** | IV | IV |
| Nedmuldning af halm før vintersæd | Fjernelse af halm før vintersæd | 0 8) | ** | IV | IV |
| Biochar | Ingen biochar produktion og tilsætning | 0 8) | IV | IV | IV |
| Positionsbestemt tilførsel af gødning | Bredspredning af handelsgødning | 1-2 kg N ha ⁻¹ | ** | IV | IV |
| Ændret udbringningsperiode for husdyrgødning om efteråret | Husdyrgødningen kan gemmes til næste forår, alternativt udbringes inden 1. september | Samlet effekt: 1850 t N | ** | 12 | 15 |
| Afbrænding af husdyrgødning | Antages kun at være aktuelt for biogasbehandlet gylle | Svinegylle: 5 kg N DE-1 Kvæggylle: 8 kg N DE-1 Fjerkrædybstr.: 20 kg N DE-1 | ** | IV | IV |
| Kontrolleret dræning | Almindelig dræning | Pt. ikke datagrundlag | * | IV | IV |
| Konstr. minivådområder med overfladisk afstrømning | Er målrettet dræntransport; virker uden for markfladen | 5-20 kg N ha ⁻¹ opland; 500-3500 kg N ha ⁻¹ anlæg | ** | 21 - 173 | 27 - 232 |
| Konstr. minivådområder med | Er målrettet dræntransport; virker | 5-35 kg N ha ⁻¹ opland; 500-7000 kg | ** | IV | IV |

| | | | | | |
|---------------------------|------------------------|--------------------------|-----|---------|--------------|
| filtermatrice | uden for markfladen | N ha-1 anlæg | | | |
| Vådområder | Jord i omdrift | 120-190 kg N ha-1 9) | *** | 31-33 | 41-44 |
| Marine virkemidler | | | | | |
| Muslingeopdræt | IR | 600-900 kg N ha-1 10) | ** | 70-97 | 93-129 11) |
| Tangdyrknings | IR | 16 kg N ha-1 10) | ** | 575-805 | 762-1068 12) |
| Udplantning af ålegræs | IR | IV | IV | IV | IV |
| Stenrev | IR | IV | IV | IV | IV |

ETTERORD

[Sett inn tekst]

| | |
|--|------------------|
| Nøkkelord: | [Sett inn tekst] |
| Key words: | [Sett inn tekst] |
| Andre aktuelle publikasjoner fra prosjekt: | [Sett inn tekst] |

UDKAS

Tabell: NIBIO venstre justert tabell – tabellstil

| asdfasd | Venstre justert | Venstre justert | Venstre justert | Venstre justert |
|-----------------|------------------------|------------------------|------------------------|------------------------|
| asdfasdf | asdfasd | asdfasd | asdfasd | asdfasdf |
| asdfasdf | asdfasdf | asdfasdf | asdfasfd | asdfasdf |
| sdfasd | asdf | asdf | asdf | asdf |

Tabell: NIBIO høyre justert tabell for tall – tabellstil

| asdfasd | Midtstilt tekst | Midtstilt tekst | Midtstilt tekst | Midtstilt tekst |
|-----------------|------------------------|------------------------|------------------------|------------------------|
| asdfasdf | 1234 | 1234 | 1234 | 1234 |
| asdfasdf | 1234 | 1234 | 1234 | 1234 |
| sdfasd | 1234 | 1234 | 1234 | 1234 |



NIBIO
NORSK INSTITUTT FOR
BIOØKONOMI

Norsk institutt for bioøkonomi (NIBIO) ble opprettet 1. juli 2015 som en fusjon av Bioforsk, Norsk institutt for landbruksøkonomisk forskning (NILF) og Norsk institutt for skog og landskap.

Bioøkonomi baserer seg på utnyttelse og forvaltning av biologiske ressurser fra jord og hav, fremfor en fossil økonomi som er basert på kull, olje og gass. NIBIO skal være nasjonalt ledende for utvikling av kunnskap om bioøkonomi.

Gjennom forskning og kunnskapsproduksjon skal instituttet bidra til matsikkerhet, bærekraftig ressursforvaltning, innovasjon og verdiskaping innenfor verdikjedene for mat, skog og andre biobaserte næringer. Instituttet skal levere forskning, forvaltningsstøtte og kunnskap til anvendelse i nasjonal beredskap, forvaltning, næringsliv og samfunnet for øvrig.

NIBIO er eid av Landbruks- og matdepartementet som et forvaltningsorgan med særskilte fullmakter og eget styre. Hovedkontoret er på Ås. Instituttet har flere regionale enheter og et avdelingskontor i Oslo.

Forsidefoto: [Sett inn fotografens navn/eventuell fjernes denne teksten]

nibio.no